

MIXED CELL DIAGNOSTIC SYSTEMS

This is a continuation-in-part application of co-pending U.S. Application Serial No. 10/407,789, filed April 4, 2003, which is a continuation of U.S. Application Serial No. 09/927,481, filed August 9, 2001, and issued on June 3, 2003 as patent No. 6,573,080, which is a continuation of U.S. Application Serial No. 09/661,849, filed September 14, 2000, and issued on April 23, 2002 as Patent No. 6,376,172, which is a divisional of U.S. Application Serial No. 09/066,072, filed April 24, 1998, which issued on January 2, 2001 as U.S. Patent No. 6,168,915, the contents of which are incorporated in their entirety for all purposes.

10 FIELD OF THE INVENTION

The present invention generally relates to the field of diagnostic microbiology, and more particularly, to compositions and methods for detecting and differentiating one or more viruses or other intracellular parasites present in a specimen. The present invention also provides compositions and methods to evaluate the susceptibility of a organisms to antimicrobial agents.

BACKGROUND OF THE INVENTION

Despite recent advances in methods for the detection of viruses using molecular methods, the detection and identification of these organisms in cell culture remains the "gold standard" by which most viral diseases are definitively diagnosed and the method against which newer methods are compared (*See e.g.*, Wiedbrauk and Johnston, *Manual of Clinical Virology*, Raven Press, Inc., New York, NY [1993], pp. 1-17). Cell cultures are also used for the detection and identification of other intracellular parasites, especially obligate intracellular parasites such as *Chlamydia* and *Rickettsia*.

There are two general types of cell culture methods used for virus identification. The first method uses identification of virus-induced cytopathic effect (CPE) as an endpoint for virus detection. The second method utilizes molecular methods to identify the presence of virus before CPE is evident in the infected cultures. Both methods utilize cell cultures, which may present problems for small laboratories with limited expertise in cell culturing methods, space, funding, equipment, and supplies. Depending upon the cells used, cell cultures can be difficult to maintain and often require the efforts of skilled laboratorians. In addition, cell cultures require equipment such as cell culture hoods,

inverted microscopes (for observation of cells), incubators with CO₂ lines, and other equipment not readily available in many laboratories.

CPE-Based Tests

CPE-based tests often require long incubation times, as virus-induced CPE only becomes evident after multiple rounds of viral replication and spread of virus to neighboring cells (*i.e.*, the cells are "permissive" for viral infection). Virus spread results in the destruction of the cells surrounding the cell originally infected. CPE-based tests have been traditionally conducted in tubes or flasks containing a single cell type that is adhered or anchored to the sides and/or bottom of the tube or flask. As the virus must infect a cell, replicate, and spread to neighboring cells in which the process is repeated, results can be delayed for at least 28 days. Indeed, results are often not available for 7-28 days after inoculation of the cell culture with the virus suspension (*See e.g.*, Leland, *Clinical Virology*, W.B. Saunders, Philadelphia [1996], pp. 60-65). The time necessary to establish visible CPE is dependent upon the rate of viral replication, which can vary among cell types and viruses. Thus, the amount of time needed to detect virus in a sample can greatly vary.

Pre-CPE Tests

In contrast to CPE-based tests, pre-CPE tests require only entry of the virus into a susceptible host cell and detectable expression of at least one early virus-specific antigen or nucleic acid. Detection of the virus-specific analyte or other indicator is accomplished by a number of methods (*e.g.*, labeled antibodies, the polymerase chain reaction [PCR], or the use of other reporters, such as the ELVISTM system). Expression of early viral genes has been shown to be very rapid in many virus-host cell systems *in vitro*. Thus, use of pre-CPE based virus tests significantly reduces the time required to detect and identify viruses in clinical specimens.

Pre-CPE detection of virus is often accomplished by using monolayers of adherent cells grown on 12 mm round coverslips contained in 1 dram shell vials (*i.e.*, the "shell vial" method or technique). The shell vial technique uses centrifugation of the specimen to force viral introduction into cells and enhance viral isolation. These vials are prepared by dispensing cells into sterile shell vials containing coverslips. The vial are incubated in an upright position until the cells form a monolayer on the coverslip. For shell vial inoculation, the culture medium is decanted from the vial, processed sample (*i.e.*, the clinical specimen) is added to the cell monolayer, and the vial is centrifuged at low speed,

often for one hour. After centrifugation, fresh culture medium is added to each vial. The vials are then incubated for the desired period of time. At the end of the incubation period, the coverslips are stained using an antigen detection method (*e.g.*, immunofluorescence) or the cells are evaluated via molecular diagnostic techniques.

5 In addition to viruses, shell vials are also commonly used for the detection and identification of *Chlamydia*, as other methods available for the detection and identification of these organisms are quite cumbersome, as well as time and reagent-consuming (*See e.g.*, Wiedbrauk and Johnston, *supra*, pp. 64-76).

10 The major advantage of these pre-CPE testing methods is that rapid test results are often possible. One major disadvantage to pre-CPE testing of shell vial cultures is that this type of test is feasible and cost-effective only if one or a few viral agents are sought for identification, and if a high proportion of specimens are likely to be positive (*See e.g.*, Schmidt and Emmons, "General Principles of Laboratory Diagnostic Methods for Viral, Rickettsial, and Chlamydial Infections," in Schmidt and Emmons (eds.), *Diagnostic Procedures for Viral, Rickettsial and Chlamydial Infections*, American Public Health Association, Washington, D.C., [1989], at p. 4).

Clinical Specimens

20 For example, the presence of skin vesicles in the genital area of a patient is highly suspicious for infection by herpes simplex virus (HSV). Typically, the physician will obtain a specimen from the affected region (*i.e.*, a vesicle) and order a CPE or a pre-CPE virus test on a single, HSV-susceptible cell line. These cell lines are often supplied either in tubes, shell vials, or multi-well plates (*e.g.*, microtiter plates). After inoculation of the cell line and an appropriate incubation time, confirmation of the presence of HSV in the sample can be accomplished using one or more of the many analytical methods (*e.g.*, immunofluorescence, immunoperoxidase, nucleic acid probes, or substrates for virus-induced reporter genes).

25 For detection of cytomegalovirus (CMV), shell vials containing cells from a single cell line (*e.g.*, human fibroblast cell lines, such as lung [MRC-5 cells] or foreskin [HFF] cells) are often used. The cells are grown to confluency on the coverslip within the vial, the sample is added to the vial, the vial is incubated for 24-48 hours or longer, and an immunofluorescent method is used to detect expression of CMV early antigen.

30 Accurate differential diagnosis is significantly more difficult in virus diseases due to respiratory, gastrointestinal, genital, or parenteral routes of transmission because many

pathogenic viruses are capable of eliciting similar symptoms or the infection is sub-clinical (*i.e.*, the signs and symptoms are not readily apparent).

Of the respiratory viruses, rhinoviruses and corona viruses are responsible for a large proportion of upper respiratory infections. Once these viruses reach the upper
5 respiratory mucosa, they attach to and infect epithelial cells. Typically, these infections last only a few days and self-resolve. Other respiratory viruses, such as the influenzas, parainfluenzas, respiratory syncytial virus (RSV), and various adenoviruses attach to and infect ciliated, columnar epithelial cells. The virus-infected cells lyse, resulting in the release of enzymes and activate complement, resulting in a local mononuclear
10 inflammatory response. Normal airway clearance mechanisms fail because of the failure of the epithelial cells to function normally. These cells may also slough off. Cell debris from dead and dying cells often obstructs airways, and the host becomes very susceptible to secondary bacterial infection and/or superinfection. All of these viruses may progress to lower respiratory involvement and pneumonia. After replication in the respiratory
15 epithelial cells, adenovirus may travel via the blood to the lymphoid tissues in all areas of the body, causing systemic infection or disease.

Standard clinical virology practice is to inoculate multiple tubes of cell cultures with the specimen (*e.g.*, throat swab, nasopharyngeal swab, or sputum specimen) as the tropism of each type of virus for specific cell types is often very narrow (*i.e.*, only one
20 type of virus may grow optimally on a single cell type). This narrow tropism of virus for a limited number of cell types creates at least two major practical problems for both CPE and pre-CPE virus testing.

First, primary monkey kidney cells are currently the cell line of choice for isolation of influenza viruses. The manufacture of these cells requires the quarantine of
25 source animals for long periods prior to sacrifice and cell culture preparation. This quarantine period is used to monitor the animals for good health and allows time to test the animals for infection by endogenous simian viruses such as foamy virus, SV5, and SV40. The quarantine period also greatly reduces, but does not eliminate, the possibility that the monkeys are infected with Monkey B Virus, a herpesvirus that is highly fatal to
30 humans. In addition, there are other problems related to the use of monkeys for the production of primary cell cultures, including the reduction in the stock of suitable animals due to importation concerns and monkey population considerations.

Second, additional continuous cell lines are required in order to detect respiratory viruses other than influenza virus. Thus, multiple cell lines are used in order to diagnose
35 the viral infection/disease of each patient. The need for multiple units of individual cell

lines is compounded in methods using pre-CPE tests for detection and identification of respiratory viruses. Pre-CPE testing for respiratory viruses requires the expenditure of significant labor in handling coverslips, the added expense of molecular reagents used with multiple cell lines for both positive and negative specimens, and the significant labor associated with microscopically reading each of the multiple cell lines inoculated in the panel of cell lines.

However, despite these drawbacks, shell vial technology using single cell types in multiple units (tubes, shell vials, etc.), is still currently used to detect respiratory viruses, as it is a proven method. For example, detection of RSV in 16 hours using shell vials containing only HEp-2 cells yielded more positives than antigen detection methods applied directly to the clinical specimen, and as many positives as conventional cell cultures (Smith *et al.*, J. Clin. Microbiol., 29:463-465 [1991]). Isolation of other respiratory viruses has also been possible with shell vial cultures containing a monolayer of a single cell type. For example, using vials of primary monkey kidney cells and A549 cells incubated for 40 hours, 83% of adenoviruses, 94% of influenza B, and 80% of parainfluenza virus types 1, 2, and 3 were identified (Rabalais *et al.*, J. Clin. Microbiol., 30:1505-1508 [1992]). In another report, 50% of adenoviruses, 94% of influenza A viruses, 100% of influenza B viruses, and 100% of parainfluenza viruses, in shell vials of primary rhesus monkey kidney cells, and 92% of RSV in shell vials of HEp-2 cells incubated for 2-4 days (*See e.g.*, Olsen *et al.*, J. Clin. Microbiol., 31:422-425 [1993]; and Leland, *Clinical Virology*, W.B. Saunders Company, Philadelphia, PA [1996], at p. 85-86).

Although these methods provide relatively rapid results (*i.e.*, as opposed to the long incubation periods often necessary for CPE tests), there remains a need in clinical and reference virology laboratories for cell culture methods and compositions for the reliable detection and identification of viruses in a single, easy-to-manipulate unit that provides rapid detection and identification in a cost-effective manner, while also providing the sensitivity of a diagnostic assay system.

SUMMARY OF THE INVENTION

The present invention generally relates to the field of diagnostic microbiology, and more particularly, to compositions and methods for detecting and differentiating one or more viruses or other intracellular parasites present in a specimen. The present invention also provides compositions and methods to evaluate the susceptibility of a organisms to antimicrobial agents.

In particular, the present invention provides methods and compositions suitable for the detection of viruses using CPE-based and pre-CPE methods. The preferred embodiments encompass mixed cell cultures which contain at least two different cell types. In some preferred embodiments, the mixed cell cultures contain two different cell types, while in other embodiments, the mixed cell cultures contain three or more different cell types. Thus, it is intended that the present invention encompass compositions in which at least two cell types are mixed together in one container (*e.g.*, flask, tube, shell vial, or any other container suitable for the growth of cells). Importantly, each cell type within these mixed cell cultures retains its susceptibility to viruses and other intracellular parasites as if it was in a single cell culture (*i.e.*, a cell culture that contains only one cell type, as known in the art). In addition, the mixed cell cultures of the present invention remain viable for as long as required for their use in diagnostic assays. In particularly preferred embodiments, the cell types included within mixed cell cultures are present in approximately the same ratio (*i.e.*, for a two cell type mixed, there is a 50:50 ratio of cell types). However, it is not intended that the present invention be limited to any particular ratio of cell types in mixed culture, as various detection systems may be optimized using different ratios. For example, in some circumstances, a two cell mixture of 45:55, 40:60, or even 35:75, may be more suited than a 50:50 ratio.

The present invention also provides methods and compositions suitable for the detection and identification of non-viral obligate intracellular and intracellular parasites, such as members of the *Chlamydiales* and *Rickettsiales*.

The present invention also contemplates compositions comprising a cell culture suitable for the detection of intracellular parasites, wherein the cell culture comprises at least two cell types susceptible to infection by at least one intracellular parasite. In some preferred embodiments of the composition, the cell types comprise a first cell type and a second cell type. In some embodiments, the first cell type consists of buffalo green monkey kidney cells and the second cell type consists of mink lung cells. In other embodiments, the first cell type consists of mink lung cells and the second cell type consists of human mucoepidermoid cells. In yet other embodiments, the first cell type consists of human lung carcinoma cells and the second cell type consists of human mucoepidermoid cells. In still other embodiments, the first cell type consists of buffalo green monkey kidney cells and the second cell type consists of human embryonic lung cells. In further embodiments, the cell type consists of human epidermoid laryngeal carcinoma cells and the second cell type consists of McCoy cells. In additional

embodiments, the first cell type consists of mink lung cells and the second cell type consists of human diploid lung cells.

5 In some preferred embodiments, the cell types of the composition are susceptible to respiratory viruses, including but not limited to influenza viruses of any type (*e.g.*, Influenza A, Influenza B, and Influenza C) and/or strain, RSV, cytomegalovirus, parainfluenza viruses, respiratory syncytial virus, rhinoviruses, coronaviruses, and adenoviruses. In yet other embodiments, the cell types of the composition are susceptible to enteroviruses, including but not limited to any type and/or strain of echovirus, poliovirus, and Coxsackie virus (*e.g.*, Coxsackie A and B viruses), and numbered EV
10 strains. In addition to enteroviruses, it is contemplated that the present invention encompass cell types that are susceptible to picornaviruses such as Hepatitis A.

The present invention also provides methods for the detection and identification of intracellular parasites in a sample, comprising the steps of providing a sample suspected of containing one or more intracellular parasites; and a mixed cell culture
15 comprising at least two cell types; inoculating the mixed cell culture with the sample to produce an inoculated culture; and observing the inoculated culture for the presence of the one or more intracellular parasites.

In some embodiments of the method, the intracellular parasite is a virus. In some particularly preferred embodiments, the virus is selected from the group consisting of
20 cytomegalovirus, influenza viruses, parainfluenza viruses, respiratory syncytial virus, rhinoviruses, coronaviruses, and adenoviruses. In yet other embodiments of the methods, the virus is an enterovirus. In other particularly preferred embodiments, the enterovirus is selected from the group consisting of poliovirus, Coxsackie viruses and echoviruses (*e.g.*, Coxsackie A and B viruses), and numbered EV strains. In addition to enteroviruses, it is
25 contemplated that the present invention encompass cell types that are susceptible to picornaviruses such as Hepatitis A. In still other preferred embodiments, the virus is a herpes virus. In other particularly preferred embodiments, the herpes virus is selected from the group consisting of Herpes Simplex Type 1, Herpes Simplex Type 2, Cytomegalovirus, Varicella-Zoster virus, Epstein-Barr virus, Human Herpes Virus 6,
30 Human Herpes Virus 7, and Human Herpes Virus 8. In yet other preferred embodiments, the intracellular parasite is a member of the genus *Chlamydia*. In still other particularly preferred embodiments, the intracellular parasite is *C. trachomatis*.

In some preferred embodiments of the methods, the cell types comprise a first cell type and a second cell type. In some preferred embodiments, the first cell type is a mink
35 lung cell, and the second cell type is a human mucoepidermoid cell. In other preferred

embodiments, the first cell type is a buffalo green monkey kidney cell and the second cell type is a human mucoepidermoid cell. In yet another alternative embodiment, the first cell type is a genetically engineered baby hamster kidney cell and the second cell type is a mink lung cell. In still other embodiments, the first cell type is a first genetically engineered cell type and the second cell type is a second genetically engineered cell type.

It is contemplated that the methods of the present invention will be used in conjunction with controls of known positivity and negativity for the virus(es) and/or other intracellular organism of interest.

The present invention also provides methods for the detection and identification of intracellular parasites in a sample, comprising the steps of providing: a sample suspected of containing one or more intracellular parasites; and a mixed cell culture comprising a first cell type and a second cell type; inoculating the mixed cell culture with the sample to produce an inoculated culture; and observing the inoculated culture for the presence of the one or more intracellular parasites.

In some particularly preferred embodiments, the intracellular parasite is a virus. In some particularly preferred embodiments, the virus is selected from the group consisting of cytomegalovirus, influenza viruses, parainfluenza viruses, respiratory syncytial virus, rhinoviruses, coronaviruses, and adenoviruses. In yet other embodiments of the methods, the virus is an enterovirus. In other particularly preferred embodiments, the enterovirus is selected from the group consisting of poliovirus, Coxsackie viruses and echoviruses (e.g., Coxsackie A and B viruses), and numbered EV strains. In addition to enteroviruses, it is contemplated that the present invention encompass cell types that are susceptible to picornaviruses such as Hepatitis A. In still other preferred embodiments, the virus is a herpes virus. In other particularly preferred embodiments, the herpes virus is selected from the group consisting of Herpes Simplex Type 1, Herpes Simplex Type 2, Cytomegalovirus, Varicella-Zoster virus, Epstein-Barr virus, Human Herpes Virus 6, Human Herpes Virus 7, and Human Herpes Virus 8. In yet other preferred embodiments, the intracellular parasite is a member of the genus *Chlamydia*. In still other particularly preferred embodiments, the intracellular parasite is *C. trachomatis*.

In some preferred embodiments of the methods, the cell types comprise a first cell type and a second cell type. In some preferred embodiments, the first cell type is a mink lung cell, and the second cell type is a human mucoepidermoid cell. In other preferred embodiments, the first cell type is a buffalo green monkey kidney cell and the second cell type is a human mucoepidermoid cell. In yet another alternative embodiment, the first cell type is a genetically engineered baby hamster kidney cell and the second cell type is a

mink lung cell. In still other embodiments, the first cell type is a first genetically engineered cell type and the second cell type is a second genetically engineered cell type.

It is contemplated that the methods of the present invention will be used in conjunction with controls of known positivity and negativity for the virus(es) and/or other intracellular organism of interest.

The present invention further provides methods for the detection of influenza virus, comprising the steps of providing a sample suspected of containing influenza virus, and mink lung cells; inoculating the mink lung cells with the sample; and detecting the presence of the influenza within the mink lung cells. In particularly preferred embodiments, the mink lung cells are Mv1Lu cells. In alternative embodiments, the influenza virus is selected from the group consisting of Influenza A, Influenza B, and Influenza C.

It is contemplated that the methods of the present invention will be used in conjunction with controls of known positivity and negativity for the virus(es) and/or other intracellular organism of interest.

In one embodiment, the present invention provides methods for the detection of infectious virus in a specimen comprising the steps of: a) providing a specimen suspected of containing a virus, a cell line permissive for infection by the virus, and a genetically engineered cell line containing an oligonucleotide having a sequence comprising a promoter sequence derived from the virus, wherein the promoter sequence is operably linked to a reporter gene, and wherein the expression of the reporter gene is dependent upon and quantitatively proportional to the presence of the virus; b) mixing together the permissive cell line and the genetically engineered cell line to create a mixed cell culture; c) inoculating the mixed cell culture with the specimen under conditions which permit the infection of the mixed cell culture by the virus; and d) detecting the expression of the reporter gene and thereby detecting the presence of virus in the specimen. In one preferred embodiment, the mixed cell culture is a mixture consisting of 80-99% of the permissive cell line and 1-20% of the genetically engineered cell line. In other preferred embodiments, the mixed cell culture is a mixture consisting of equal proportions of the cell types used in the mixture.

In one embodiment of the method, the genetically engineered cell line contains an oligonucleotide having a sequence comprising a herpesvirus inducible promoter operably linked to a reporter gene selected from the group comprising the *Escherichia coli lacZ* gene and a luciferase gene. In one preferred embodiment of the method, the genetically engineered cell line is BHKICP10LacZ. In an alternative preferred embodiment, the

genetically engineered cell line is BHKICP6LacZ. However, it is not intended that the reporter gene be limited to the *lacZ* and luciferase genes. Indeed, it is contemplated that any suitable reporter gene known to those in the art will be useful in the method of the present invention.

5 It is also contemplated that various permissive cell lines will be useful in the method of the present invention. In one embodiment, the permissive cell line is permissive for infection with herpesvirus. In a particularly preferred embodiment, the permissive cell line is MRC-5.

10 It is contemplated that the method of the present invention will be used in conjunction with controls of known positivity and negativity for the virus(es) of interest. Thus, for mixed cultures in which genetically engineered cell lines are used, it is contemplated that the pattern of reporter gene expression present in a test sample (*e.g.*, from a clinical specimen) will be compared to the patterns of reporter gene expression in control samples known to be positive and/or negative for the virus(es) of interest. It is
15 also contemplated that effects unrelated to the expression of the reporter gene will be detectable, including but not limited to CPE. These effects, alone and in combination with the reporter gene expression may be used to detect the presence of viral infection.

 The present invention also provides methods for the typing of infectious herpesvirus in specimens, comprising the steps of: a) providing a specimen suspected of
20 containing one or more members of the herpesvirus family, a cell line permissive for infection by one or more members of the herpesvirus family, a genetically engineered cell line containing an oligonucleotide having a sequence comprising a promoter sequence derived from a member of the herpesvirus family wherein the promoter sequence is operably linked to a reporter gene, and the expression of the reporter gene is dependent
25 upon and quantitatively proportional to the presence of herpesvirus and wherein the expression of the reporter gene varies in a distinguishable manner as a result of the presence of different members of the herpesvirus family; b) mixing together the permissive cell line and the genetically engineered cell line to create a mixed cell culture; c) inoculating this mixed cell culture with the specimen under conditions which permit
30 the infection of the mixed cell culture by members of the herpesvirus family, wherein the infection results in a distinguishable pattern of expression by the reporter gene; d) detecting the expression of the reporter gene and thereby detecting the presence of one or more members of the herpesvirus family in the specimen; and e) identifying the presence of a specific member of the herpesvirus family based upon the resulting distinguishable
35 pattern. It is contemplated that this pattern of expression will be observable by various

assisted and non-assisted methods, including visual observation by eye, spectrophotometric observation, etc. It is not intended that the detection of distinguishable pattern(s) be limited to any particular method of detection.

5 In a preferred embodiment of the typing method of the present invention, the mixed cell culture is a mixture consisting of 80-99% of the permissive cell line and 1-20% of the genetically engineered cell line. In yet other preferred embodiments, the cell types are in approximate equal proportions in the mixed cell cultures. As with the first method described, it is not intended that the present invention be limited to any particular herpesvirus. In one particular embodiment, the member of the herpesvirus
10 family detected and typed using the method of the present invention is selected from the group comprising HSV-1, HSV-2, CMV, VZV, EBV, and human herpes viruses such as HHV-6, HHV-7, and HHV-8. It is intended that one or more herpesviruses may be detected and typed in one specimen. In this manner, co-infection with multiple herpesviruses may be diagnosed. For example, it is contemplated that mixed infections
15 with HSV-1 and HSV-2 may be detectable and the infections distinguished using the methods of the present invention.

In one embodiment of the typing method, the reporter gene comprises *E. coli lacZ* gene. However, it is not intended that the reporter gene be limited to *lacZ*. Indeed, it is contemplated that any reporter gene may be used in this method. In one particularly
20 preferred embodiment, the detection of the reporter gene is accomplished through the use of histochemical staining. It is contemplated that one member of the herpesvirus family will produce an histochemically pattern of expression that is distinguishable from the histochemical patterns produced by other members of the herpesvirus family. In this manner, it is possible to use the methods of the present invention to distinguish infection
25 with one herpesvirus from infection with another herpesvirus.

It is contemplated that the method of the present invention will be used in conjunction with controls of known positivity and negativity for the virus(es) of interest. Thus, it is contemplated that the pattern of expression present in a test sample (*e.g.*, from a clinical specimen) will be compared to the patterns of expression in control samples
30 known to be positive and/or negative for the virus(es) of interest. It is also contemplated that effects unrelated to the expression of the reporter gene will be detectable, including but not limited to CPE. These effects, alone and in combination with reporter gene expression may be used to detect the presence of viral infection, as well as provide information to distinguish between viruses.

In yet another embodiment, the present invention provides a composition comprising a mixed cell culture, wherein the mixed cell culture comprises the combination of a genetically engineered cell line transformed with a promoter sequence from a virus, wherein the promoter sequence is operably linked to a reporter gene, and wherein expression of the reporter gene is dependent upon and quantitatively proportional to the presence of virus, and a non-engineered cell line which is permissive for virus infection.

In one embodiment of the composition, the mixed cell culture is a mixture consisting of 1-20% of the genetically engineered cell line and 80-99% of the permissive cell line. In yet other preferred embodiments, the cell types are in approximate equal proportions in the mixed cell cultures. In one preferred embodiment of the composition, the genetically engineered cell line component may comprise a promoter for a gene that encodes ribonucleotide reductase. In an alternative preferred embodiment, the promoter may comprise genes that encode one or more subunits of ribonucleotide reductase. In one particularly preferred embodiment, the genetically engineered cell line is BHKICP10LacZ, while in another particularly preferred embodiment, the genetically engineered cell line is BHKICP6LacZ. In an alternative embodiment of the composition, the genetically engineered cell line comprises an *E. coli lacZ* gene positioned 3' to a virus inducible promoter. It is contemplated that this *lacZ* gene be positioned immediately 3' to this virus-inducible promoter. However, it is not intended that these sequences will be contiguous. Indeed, it is contemplated only that the reporter and promoter genes are operably linked. Furthermore, it is contemplated that the composition will comprise promoter sequences from any virus, including but not limited to members of the herpesvirus family. It is also contemplated that the non-engineered cell line be permissive for infection by any number of viruses, including but not limited to members of the herpesvirus family.

In one preferred embodiment, the composition includes a genetically engineered cell line which includes a promoter for a gene that encodes a ribonucleotide reductase large subunit and the virus is a member of the herpesvirus family selected from the group consisting of HSV-1, HSV-2, CMV, VZV, EBV, HHV-6, HHV-7, and HHV-8. However, it is not intended that the present invention be limited to any particular herpesvirus. In one preferred embodiment, the genetically engineered cell line component contains an ICP10 promoter and the herpesvirus family member is HSV-2, while in another preferred embodiment, the genetically engineered cell line comprises an ICP6 promoter and the herpesvirus family member is HSV-1.

It is contemplated that the detection of reporter gene expression be accomplished through the use of various methods, including, but not limited to colorimetric, fluorimetric or luminometric assays or assay systems. In one preferred embodiment, the reporter gene encodes β -galactosidase.

5 In one embodiment, the composition includes a genetically engineered cell line that is a mammalian cell line susceptible to infection by virus. In one preferred embodiment, the genetically engineered cell line comprises baby hamster kidney cells (e.g., cell lines derived from BHK cells). In one embodiment, the composition includes a permissive cell line that is permissive to infection by herpesviruses, including but not
10 limited to HSV-1 and HSV-2. In a particularly preferred embodiment, the permissive cell line is MRC-5. It is not intended that the composition of the present invention be limited to detection of viral infection based on the expression of the reporter gene, as effects such as CPE may also be detectable.

The present invention also provides a kit for assaying for the presence of
15 infectious herpesvirus in a specimen. The kit includes: a) a supply of a mixed cell line comprised of a cell line of genetically engineered mammalian cells susceptible to infection by herpesvirus, wherein the cell line contains an oligonucleotide having a sequence comprising a virus promoter sequence operably linked to a reporter gene, and where the expression of the reporter gene is dependent upon and quantitatively
20 proportional to the presence of virus in the specimen; and a cell line permissive for virus; and b) a supply of reagents to detect the expression of the reporter gene. It is not intended that the promoter sequences present within the genetically engineered cell line be limited to any particular virus or virus family. It is contemplated that any virus promoter will be useful in the kit of the present invention. However, in one preferred embodiment,
25 herpesvirus promoter sequences are present in the genetically engineered cell line.

It is contemplated that various promoter sequences will be useful in the kit of the present invention. However, in a preferred embodiment, the promoter encodes either a complete ribonucleotide reductase enzyme, or in the alternative, subunits of
30 ribonucleotide reductase. In one particularly preferred embodiment, the promoter sequence contains a promoter for a gene that encodes a ribonucleotide reductase large subunit and the herpesvirus is a herpesvirus family member selected from the group consisting of HSV-1, HSV-2, CMV, VZV, EBV, HHV-6, HHV-7, and HHV-8. However, it is not intended that the kit will be limited to this list of herpesviruses. Indeed, it is contemplated that any herpesvirus may be detected using the present kit. In
35 one particularly preferred embodiment of the kit, the promoter sequence contains an

ICP10 promoter and the herpesvirus family member is HSV-2, while in an alternative preferred embodiment, the promoter sequence contains an ICP6 promoter and the herpesvirus family member is HSV-1.

5 In one preferred embodiment of the kit, the promoter sequence present in the genetically engineered cell line comprises an *E. coli lacZ* gene that is operably linked to a herpesvirus inducible promoter. In one particularly preferred embodiment, the genetically engineered mammalian cells are BHKICP10LacZ cells, while in an alternative embodiment, the cells are BHKICP6LacZ cells.

10 In one preferred embodiment, the reporter gene encodes β -galactosidase. However, it is not intended that the present invention be limited to any particular reporter gene. It is also contemplated that the reporter gene will encode any number of enzymes that are amenable to detection by various methods, including but not limited to such methods as colorimetric, fluorimetric or luminometric assay systems. In one preferred embodiment of the kit, the reagents provided for the detection of reporter gene expression
15 may include, but are not limited to, solutions of 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside, o-nitrophenyl-galactopyranoside solution, and fluorescein di- β -D-galactopyranoside. However, it is not intended to limit the kit to these assay systems, as other systems (*e.g.*, radiometric assay systems) may be useful.

20 It is contemplated that the kit of the present invention may also include additional components, such as materials suitable for positive and negative controls and instructions for use. It is not intended that the kit of the present invention be limited to the mixed cell line and reagents for the detection of reporter gene expression. It is also intended that the kit will be useful for detection of viral effects on cells other than and unrelated to reporter gene expression. For example, it is contemplated that the kit may be useful for detection
25 of CPE.

30 In a further embodiment, the invention provides a composition comprising a mixed cell culture comprising MDCK cells and one or more of A549 cells and H292 cells. These compositions are useful in detecting the presence of one or more of influenza viruses (such as influenza A and/or B), respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus. These methods are also useful in producing one or more of influenza
35 viruses (such as influenza A and/or B), respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, and parainfluenza 3 virus.

Also provided herein is a method for detecting influenza virus, comprising: 1)
35 providing: a) mixed cell culture comprising MDCK cells and one or more of A549 cells

and H292 cells; and b) a sample; 2) inoculating the mixed cell culture with the sample to produce an inoculated culture; and 3) detecting the presence of influenza virus. In one embodiment, the level of permissivity and/or susceptibility of the MDCK cells to severe acute respiratory syndrome coronavirus is 0.004% the level of permissivity and/or susceptibility of Mv1Lu cells to severe acute respiratory syndrome coronavirus. In a further embodiment, the influenza virus comprises one or more of influenza A virus and influenza B virus. In another embodiment, the mixed cell culture comprises MDCK cells and A549 cells, and the method optionally further comprises detecting the presence of one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus. In an alternative embodiment, the mixed cell culture comprises MDCK cells and H292 cells, and the method optionally further comprises detecting the presence of one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus. In a further embodiment, the mixed cell culture comprises MDCK cells, A549 cells, and H292 cells, and the method optionally further comprises detecting the presence of one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus.

Also provided by the invention is a method for producing influenza virus, comprising: 1) providing: a) mixed cell culture comprising MDCK cells and one or more of A549 cells and H292 cells; and b) a sample; 2) inoculating the mixed cell culture with the sample to produce an inoculated culture, wherein the inoculated culture produces influenza virus. In one embodiment, the influenza virus comprises one or more of influenza A virus and influenza B virus. In another embodiment, the mixed cell culture comprises MDCK cells and A549 cells, and the method optionally further comprises producing one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus. Alternatively, the mixed cell culture comprises MDCK cells and H292 cells, and the method optionally further comprises producing one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus. In yet another alternative, the mixed cell culture comprises MDCK cells, A549 cells, and H292 cells, and the method further comprises producing one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus.

Also provided is a method for detecting metapneumovirus, comprising: 1) providing a) a mixed cell culture comprising MDCK cells and A549 cells; and b) sample; 2) inoculating the mixed cell culture with the sample to produce inoculated cells; and 3) detecting the presence of metapneumovirus. In one embodiment, the method further comprises detecting influenza virus, as exemplified by influenza B virus and/or influenza A virus. In another embodiment, the method further comprises detecting the presence of one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, and parainfluenza 3 virus. In an alternative embodiment, the mixed cell culture further comprises H292 cells, and the method optionally further comprises detecting the presence of one or more of respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, and parainfluenza 3 virus.

The invention also provides a method for producing metapneumovirus, comprising: 1) providing a mixed cell culture comprising MDCK cells and A549 cells; and b) sample; 2) inoculating the cultured cells with the sample to produce inoculated an inoculated culture, wherein the inoculated culture produces metapneumovirus. In one preferred embodiment, the mixed cell culture further comprises H292 cells.

Definitions

To facilitate understanding of the invention, a number of terms are defined below.

The terms "sample" and "specimen" in the present specification and claims are used in their broadest sense. On the one hand, they are meant to include a specimen or culture. On the other hand, they are meant to include both biological and environmental samples. These terms encompasses all types of samples obtained from humans and other animals, including but not limited to, body fluids such as urine, blood, fecal matter, cerebrospinal fluid (CSF), semen, sputum, and saliva, as well as solid tissue. These terms also refers to swabs and other sampling devices which are commonly used to obtain samples for culture of microorganisms.

Biological samples may be animal, including human, fluid or tissue, food products and ingredients such as dairy items, vegetables, meat and meat by-products, and waste. Environmental samples include environmental material such as surface matter, soil, water, and industrial samples, as well as samples obtained from food and dairy processing instruments, apparatus, equipment, disposable, and non-disposable items. These examples are not to be construed as limiting the sample types applicable to the present invention.

Whether biological or environmental, a sample suspected of containing microorganisms may (or may not) first be subjected to an enrichment means to create a "pure culture" of microorganisms. By "enrichment means" or "enrichment treatment," the present invention contemplates (i) conventional techniques for isolating a particular microorganism of interest away from other microorganisms by means of any culture medium and/or technique, and (ii) novel techniques for isolating particular microorganisms away from other microorganisms. It is not intended that the present invention be limited only to one enrichment step or type of enrichment means. For example, it is within the scope of the present invention, following subjecting a sample to a conventional enrichment means, to subject the resultant preparation to further purification such that a pure culture of a strain of a species of interest is produced. This pure culture may then be analyzed by the medium and method of the present invention.

As used herein, the term "organism" and "microorganism," are used to refer to any species or type of microorganism, including but not limited to viruses and bacteria, including rickettsia and chlamydia. Thus, the term encompasses, but is not limited to DNA and RNA viruses, as well as organisms within the orders *Rickettsiales* and *Chlamydiales*.

As used herein, the term "culture," refers to any sample or specimen which is suspected of containing one or more microorganisms. "Pure cultures" are cultures in which the organisms present are only of one strain of a particular genus and species. This is in contrast to "mixed cultures," which are cultures in which more than one genus and/or species of microorganism are present.

As used herein, the term "cell type," refers to any cell, regardless of its source or characteristics.

As used herein, the term "cell line," refers to cells that are cultured *in vitro*, including primary cell lines, finite cell lines, continuous cell lines, and transformed cell lines.

As used herein, the terms "primary cell culture," and "primary culture," refer to cell cultures that have been directly obtained from animal or insect tissue. These cultures may be derived from adults as well as fetal tissue.

As used herein, the term "finite cell lines," refer to cell cultures that are capable of a limited number of population doublings prior to senescence.

As used herein, the term "continuous cell lines," refer to cell cultures that have undergone a "crisis" phase during which a population of cells in a primary or finite cell line apparently ceases to grow, but yet a population of cells emerges with the general

characteristics of a reduced cell size, higher growth rate, higher cloning efficiency, increased tumorigenicity, and a variable chromosomal complement. These cells often result from spontaneous transformation *in vitro*. These cells have an indefinite lifespan.

As used herein, the term "transformed cell lines," refers to cell cultures that have been transformed into continuous cell lines with the characteristics as described above. Transformed cell lines can be derived directly from tumor tissue and also by *in vitro* transformation of cells with whole virus (e.g., SV40 or EBV), or DNA fragments derived from a transforming virus using vector systems.

As used herein, the term "hybridomas," refers to cells produced by fusing two cell types together. Commonly used hybridomas include those created by the fusion of antibody-secreting B cells from an immunized animal, with a malignant myeloma cell line capable of indefinite growth *in vitro*. These cells are cloned and used to prepare monoclonal antibodies.

As used herein, the term "mixed cell culture," refers to a mixture of two types of cells. In some preferred embodiments, the cells are cell lines that are not genetically engineered, while in other preferred embodiments the cells are genetically engineered cell lines. In some embodiments, the one or more of the cell types is re "permissive" (*i.e.*, virus is capable of replication and spread from cell to cell within the culture). The present invention encompasses any combination of cell types suitable for the detection, identification, and/or quantitation of viruses in samples, including mixed cell cultures in which all of the cell types used are not genetically engineered, mixtures in which one or more of the cell types are genetically engineered and the remaining cell types are not genetically engineered, and mixtures in which all of the cell types are genetically engineered.

As used herein, the term "suitable for the detection of intracellular parasites," refers to cell cultures that can be successfully used to detect the presence of an intracellular parasite in a sample. In preferred embodiments, the cell cultures are capable of maintaining their susceptibility to infection and/or support replication of the intracellular parasite. It is not intended that the present invention be limited to a particular cell type or intracellular parasite.

As used herein, the term "susceptible to infection" refers to the ability of a cell to become infected with virus or another intracellular organism. Although it encompasses "permissive" infections, it is not intended that the term be so limited, as it is intended that the term encompass circumstances in which a cell is infected, but the organism does not necessarily replicate and/or spread from the infected cell to other cells. The phrase "viral

proliferation," as used herein describes the spread or passage of infectious virus from a permissive cell type to additional cells of either a permissive or susceptible character.

As used herein, the terms "monolayer," "monolayer culture," and "monolayer cell culture," refer to cells that have adhered to a substrate and grow in as a layer that is one cell in thickness. Monolayers may be grown in any format, including but not limited to flasks, tubes, coverslips (e.g., shell vials), roller bottles, etc. Cells may also be grown attached to microcarriers, including but not limited to beads.

As used herein, the term "suspension," and "suspension culture," refers to cells that survive and proliferate without being attached to a substrate. Suspension cultures are typically produced using hematopoietic cells, transformed cell lines, and cells from malignant tumors.

As used herein, the terms "culture media," and "cell culture media," refers to media that are suitable to support the growth of cells *in vitro* (i.e., cell cultures). It is not intended that the term be limited to any particular culture medium. For example, it is intended that the definition encompass outgrowth as well as maintenance media. Indeed, it is intended that the term encompass any culture medium suitable for the growth of the cell cultures of interest.

As used herein, the term "obligate intracellular parasite," (or "obligate intracellular organism") refers to any organism which requires an intracellular environment for its survival and/or replication. Obligate intracellular parasites include viruses, as well as many other organisms, including certain bacteria (e.g., most members of the orders *Rickettsiales* [e.g., *Coxiella*, *Rickettsia* and *Ehrlichia*] and *Chlamydiales* [e.g., *C. trachomatis*, *C. psittaci*], etc). The term "intracellular parasite," refers to any organism that may be found within the cells of a host animal, including but not limited to obligate intracellular parasites briefly described above. For example, intracellular parasites include organisms such as *Brucella*, *Listeria*, *Mycobacterium* (e.g., *M. tuberculosis* and *M. leprae*), and *Plasmodium*, as well as *Rochalimea*.

As used herein, the term "antimicrobial," is used in reference to any compound which inhibits the growth of, or kills microorganisms. It is intended that the term be used in its broadest sense, and includes, but is not limited to compounds such as antibiotics which are produced naturally or synthetically. It is also intended that the term includes compounds and elements that are useful for inhibiting the growth of, or killing microorganisms.

As used herein, the terms "chromogenic compound," and "chromogenic substrate," refer to any compound useful in detection systems by their light absorption or

emission characteristics. The term is intended to encompass any enzymatic cleavage products, soluble, as well as insoluble, which are detectable either visually or with optical machinery. Included within the designation "chromogenic" are all enzymatic substrates which produce an end product which is detectable as a color change. This includes, but is not limited to any color, as used in the traditional sense of "colors," such as indigo, blue, red, yellow, green, orange, brown, etc., as well as fluorochromic or fluorogenic compounds, which produce colors detectable with fluorescence (e.g., the yellow-green of fluorescein, the red of rhodamine, etc.). It is intended that such other indicators as dyes (e.g., pH) and luminogenic compounds be encompassed within this definition.

As used herein, the commonly used meaning of the terms "pH indicator," "redox indicator," and "oxidation-reduction indicator," are intended. Thus, "pH indicator," encompasses all compounds commonly used for detection of pH changes, including, but not limited to phenol red, neutral red, bromthymol blue, bromcresol purple, bromcresol green, bromchlorophenol blue, m-cresol purple, thymol blue, bromcresol purple, xylenol blue, methyl red, methyl orange, and cresol red. The terms "redox indicator," and "oxidation-reduction indicator," encompasses all compounds commonly used for detection of oxidation/reduction potentials (*i.e.*, "eH") including, but not limited to various types or forms of tetrazolium, resazurin, and methylene blue.

As used herein, the term "inoculating suspension," or "inoculant," is used in reference to a suspension which may be inoculated with organisms to be tested. It is not intended that the term "inoculating suspension," be limited to a particular fluid or liquid substance. For example, inoculating suspensions may be comprised of water, saline, or an aqueous solution. It is also contemplated that an inoculating suspension may include a component to which water, saline or any aqueous material is added. It is contemplated in one embodiment, that the component comprises at least one component useful for the intended microorganism. It is not intended that the present invention be limited to a particular component.

As used herein, the term "kit," is used in reference to a combination of reagents and other materials.

As used herein, the term "primary isolation," refers to the process of culturing organisms directly from a sample. As used herein, the term "isolation," refers to any cultivation of organisms, whether it be primary isolation or any subsequent cultivation, including "passage," or "transfer," of stock cultures of organisms for maintenance and/or use.

As used herein, the term "presumptive diagnosis," refers to a preliminary diagnosis which gives some guidance to the treating physician as to the etiologic organism involved in the patient's disease. Presumptive diagnoses are often based on "presumptive identifications," which as used herein refer to the preliminary identification of a microorganism.

As used herein, the term "definitive diagnosis," is used to refer to a final diagnosis in which the etiologic agent of the patient's disease has been identified. The term "definitive identification" is used in reference to the final identification of an organism to the genus and/or species level.

The term "recombinant DNA molecule," as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

DNA molecules are said to have "5' ends" and "3' ends" because mononucleotides are reacted to make oligonucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage. Therefore, an end of an oligonucleotides is referred to as the "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose ring and as the "3' end" if its 3' oxygen is not linked to a 5' phosphate of a subsequent mononucleotide pentose ring. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide, also may be said to have 5' and 3' ends. In either a linear or circular DNA molecule, discrete elements are referred to as being "upstream" or 5' of the "downstream" or 3' elements. This terminology reflects the fact that transcription proceeds in a 5' to 3' fashion along the DNA strand. The promoter and enhancer elements which direct transcription of a linked gene are generally located 5' or upstream of the coding region (enhancer elements can exert their effect even when located 3' of the promoter element and the coding region). Transcription termination and polyadenylation signals are located 3' or downstream of the coding region.

The term "an oligonucleotide having a nucleotide sequence encoding a gene," refers to a DNA sequence comprising the coding region of a gene or, in other words, the DNA sequence which encodes a gene product. The coding region may be present in either a cDNA or genomic DNA form. Suitable control elements such as enhancers, promoters, splice junctions, polyadenylation signals, etc. may be placed in close proximity to the coding region of the gene if needed to permit proper initiation of transcription and/or correct processing of the primary RNA transcript. Alternatively, the coding region utilized in the vectors of the present invention may contain endogenous

enhancers and/or promoters, splice junctions, intervening sequences, polyadenylation signals, etc. or a combination of both endogenous and exogenous control elements.

The term "transcription unit," as used herein refers to the segment of DNA between the sites of initiation and termination of transcription and the regulatory elements necessary for the efficient initiation and termination. For example, a segment of DNA comprising an enhancer/promoter, a coding region, and a termination and polyadenylation sequence comprises a transcription unit.

The term "regulatory element," as used herein refers to a genetic element which controls some aspect of the expression of nucleic acid sequences. For example, a promoter is a regulatory element which facilitates the initiation of transcription of an operably linked coding region. Other regulatory elements are splicing signals, polyadenylation signals, termination signals, etc. (defined *infra*).

The terms "reporter gene construct," or "reporter gene vector," as used herein refers to a recombinant DNA molecule containing a sequence encoding the product of a reporter gene and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular host organism. Eukaryotic cells are known to utilize promoters, enhancers, and termination and polyadenylation signals.

The term "reporter gene," refers to an oligonucleotide having a sequence encoding a gene product (typically an enzyme) which is easily and quantifiably assayed when a construct comprising the reporter gene sequence operably linked to a heterologous promoter and/or enhancer element is introduced into cells containing (or which can be made to contain) the factors necessary for the activation of the promoter and/or enhancer elements. Examples of reporter genes include but are not limited to bacterial genes encoding β -galactosidase (*lacZ*), the bacterial chloramphenicol acetyltransferase (*cat*) genes, firefly luciferase genes and genes encoding β -glucuronidase (*GUS*).

Transcriptional control signals in eukaryotes comprise "promoter" and "enhancer" elements. Promoters and enhancers consist of short arrays of DNA sequences that interact specifically with cellular proteins involved in transcription (Maniatis, *et al.*, Science 236:1237 [1987]). Promoter and enhancer elements have been isolated from a variety of eukaryotic sources including genes in yeast, insect and mammalian cells and viruses (analogous control elements [*i.e.*, promoters, are also found in prokaryotes]). The selection of a particular promoter and enhancer depends on what cell type is to be used to express the protein of interest. Some eukaryotic promoters and enhancers have a broad host range while others are functional in a limited subset of cell types (for review see Voss, *et al.*, Trends Biochem. Sci., 11:287 [1986], and Maniatis, *et al.*, *supra* [1987]).

For example, the SV40 early gene enhancer is very active in a wide variety of cell types from many mammalian species and has been widely used for the expression of proteins in mammalian cells (Dijkema, *et al.*, EMBO J. 4:761 [1985]). Two other examples of promoter/enhancer elements active in a broad range of mammalian cell types are those from the human elongation factor 1 α gene (Uetsuki *et al.*, J. Biol. Chem., 264:5791 [1989]; Kim *et al.*, Gene 91:217 [1990]; and Mizushima and Nagata, Nuc. Acids. Res., 18:5322 [1990]) and the long terminal repeats of the Rous sarcoma virus (Gorman *et al.*, Proc. Natl. Acad. Sci. USA 79:6777 [1982]), and the human cytomegalovirus (Boshart *et al.*, Cell 41:521 [1985]).

The term "promoter/enhancer," denotes a segment of DNA which contains sequences capable of providing both promoter and enhancer functions (for example, the long terminal repeats of retroviruses contain both promoter and enhancer functions). The enhancer/promoter may be "endogenous," or "exogenous," or "heterologous." An endogenous enhancer/promoter is one which is naturally linked with a given gene in the genome. An exogenous (heterologous) enhancer/promoter is one which is placed in juxtaposition to a gene by means of genetic manipulation (*i.e.*, molecular biological techniques).

The presence of "splicing signals," on an expression vector often results in higher levels of expression of the recombinant transcript. Splicing signals mediate the removal of introns from the primary RNA transcript and consist of a splice donor and acceptor site (Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, New York [1989], pp. 16.7-16.8). A commonly used splice donor and acceptor site is the splice junction from the 16S RNA of SV40.

Efficient expression of recombinant DNA sequences in eukaryotic cells requires signals directing the efficient termination and polyadenylation of the resulting transcript. Transcription termination signals are generally found downstream of the polyadenylation signal and are a few hundred nucleotides in length. The term "poly A site," or "poly A sequence," as used herein denotes a DNA sequence which directs both the termination and polyadenylation of the nascent RNA transcript. Efficient polyadenylation of the recombinant transcript is desirable as transcripts lacking a poly A tail are unstable and are rapidly degraded. The poly A signal utilized in an expression vector may be "heterologous" or "endogenous." An endogenous poly A signal is one that is found naturally at the 3' end of the coding region of a given gene in the genome. A heterologous poly A signal is one which is isolated from one gene and placed 3' of another gene. A commonly used heterologous poly A signal is the SV40 poly A signal. The SV40 poly A

signal is contained on a 237 bp *Bam*HI/*Bcl*II restriction fragment and directs both termination and polyadenylation (Sambrook, *supra*, at 16.6-16.7). This 237 bp fragment is contained within a 671 bp *Bam*HI/*Pst*I restriction fragment.

5 The term "genetically engineered cell line," refers to a cell line that contains heterologous DNA introduced into the cell line by means of molecular biological techniques (*i.e.*, recombinant DNA technology).

10 The term "stable transfection," or "stably transfected," refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant," refers to a cell which has stably integrated foreign DNA into the genomic DNA.

The term "stable transfection" (or "stably transfected"), refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant," refers to a cell which has stably integrated foreign DNA into the genomic DNA.

15 The term "selectable marker," as used herein refers to the use of a gene which encodes an enzymatic activity that confers resistance to an antibiotic or drug upon the cell in which the selectable marker is expressed. Selectable markers may be "dominant"; a dominant selectable marker encodes an enzymatic activity which can be detected in any mammalian cell line. Examples of dominant selectable markers include the bacterial aminoglycoside 3' phosphotransferase gene (also referred to as the *neo* gene) which
20 confers resistance to the drug G418 in mammalian cells, the bacterial hygromycin G phosphotransferase (*hyg*) gene which confers resistance to the antibiotic hygromycin and the bacterial xanthine-guanine phosphoribosyl transferase gene (also referred to as the *gpt* gene) which confers the ability to grow in the presence of mycophenolic acid. Other
25 selectable markers are not dominant in that their use must be in conjunction with a cell line that lacks the relevant enzyme activity. Examples of non-dominant selectable markers include the thymidine kinase (*tk*) gene which is used in conjunction with *tk* cell lines, the CAD gene which is used in conjunction with CAD-deficient cells and the mammalian hypoxanthine-guanine phosphoribosyl transferase (*hprt*) gene which is used
30 in conjunction with *hprt* cell lines. A review of the use of selectable markers in mammalian cell lines is provided in Sambrook *et al.*, *supra* at pp.16.9-16.15.

The terms "nucleic acid molecule encoding," "DNA sequence encoding," and "DNA encoding," refer to the order or sequence of deoxyribonucleotides along a strand of deoxyribonucleic acid. The order of these deoxyribonucleotides determines the order of

amino acids along the polypeptide (protein) chain. The DNA sequence thus codes for the amino acid sequence.

The terms "confluent" or "confluency" as used herein in reference to an adherent cell line define a condition wherein cells throughout a culture are in contact with each other creating what appears to be a continuous sheet or "monolayer" of cells.

The terms "cytopathic effect" or "CPE" as used herein describe changes in cellular structure (*i.e.*, a pathologic effect) resulting from external agents such viruses. Common cytopathic effects include cell destruction, syncytia (*i.e.*, fused giant cells) formation, cell rounding vacuole formation, and formation of inclusion bodies. CPE results from actions of a virus on permissive cells that negatively affect the ability of the permissive cellular host to preform its required functions to remain viable. In *in vitro* cell culture systems, CPE is evident when cells, as part of a confluent monolayer, show regions of non-confluence after contact with a specimen that contains a virus. The observed microscopic effect is generally focal in nature and the foci is initiated by a single virion. However, depending upon viral load in the sample, CPE may be observed throughout the monolayer after a sufficient period of incubation. Cells demonstrating viral induced CPE usually change morphology to a rounded shape, and over a prolonged period of time can die and be released form their anchorage points in the monolayer. When many cells reach the point of focal destruction, the area is called a viral plaque, which appears as a hole in the monolayer. Cytopathic effects are readily discernable and distinguishable by those skilled in the art.

The abbreviation "ONPG," represents o-Nitrophenyl- β -D-Galactopyranoside. ONPG is a substrate for the enzyme β -galactosidase (β -gal). The reaction between ONPG and β -gal produces a yellow product which can be quantified spectrophotometrically at 405 nm.

The abbreviation "X-gal," represents the chemical compound 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside, a substrate for the enzyme β -galactosidase. The reaction between X-gal and β -galactosidase results in the formation of a blue precipitate which is visually discernable.

The term "hybriwix," represents a product of Diagnostic Hybrids, Inc., Athens, Ohio which allows for quantification of certain viral DNA in an infected monolayer of cells by DNA hybridization. "DNA hybridization" is the annealing of two complementary DNA molecules whose base sequences match according to the rules of base pairing. DNA hybridization is used to identify or quantify an unknown or "target" DNA by hybridization to a known DNA or "probe." The probe is typically labeled with a

reporter molecule such as ^{125}I , a radioisotope which can be detected and quantified with a gamma counter.

5 The phrase "plaque reduction assay," or "PRA," as used herein describes a standard method used to determine efficacy of anti-viral drugs by enumerating a decrease in plaque formation in a cell monolayer exposed to a drug. A "plaque" is a defined area of "CPE." It is usually the result of infection of the cell monolayer with a single infectious virus which then replicates and spreads to adjacent cells of the monolayer. A plaque may also be referred to as a "focus of viral infection."

10 The term "permissive" as used herein describes the sequence of interactive events between a virus and its putative host cell. The process begins with viral adsorption to the host cell surface and ends with release of infectious virions. A cell is "permissive" if it readily permits the spread of virus to other cells. Many methods are available for the determination of the permissiveness of a given cell line, including, but not limited to plaque reduction assays, comparisons of the production and/or quantitation of viral
15 proteins based on results obtained from gel electrophoresis, relative comparisons using hybridization analysis to analyze DNA or RNA content, etc.

The term "susceptible," as used herein describes the extent that a permissive or non-permissive host cell can adsorb and be penetrated by a virus. A cell line may be susceptible without being permissive in that it can be penetrated but not release virions.
20 A permissive cell line however must be susceptible.

The phrase "seed on," as used herein describes the act of transferring an aqueous solution of suspended cells into a vessel containing cells adhered to a surface, after which the vessel is stored for a sufficient period of time to allow the suspended cells or "seeds" to settle out by gravity and attach in a relatively uniform manner to the adhered cells and
25 become integrated into the final cell monolayer as a mixture. A "mixed cell monolayer," results from the "seed on" process.

The phrase "seed in," as used herein describes the mixing of two or more aqueous solutions of suspended tissue culture cells, each cell suspension having different cellular properties, and transfer of such mixture of cells into a vessel which is stored for a
30 sufficient period of time to allow the suspended cells to settle out by gravity and attach in a relatively uniform manner such that the distribution of any single cell type is indicative of the relative ratio of the cells in the original mixture.

The term "starts," as used herein refers to the reporter cells which represent a primary infection of virus. The virus infects a reporter cell (a genetically engineered cell)

and induces the expression of the reporter gene. A reporter cell can be non-permissive (*i.e.* permissiveness of the reporter cells is not required) and still produce starts.

DESCRIPTION OF THE INVENTION

5 The present invention generally relates to the field of diagnostic microbiology, and more particularly, to compositions and methods for detecting and differentiating one or more viruses or other intracellular parasites present in a specimen. The present invention also provides compositions and methods to evaluate the susceptibility of a organisms to antimicrobial agents.

10 The present invention provides methods and compositions for the detection of several different viruses, as well as other intracellular organisms present in clinical and other specimens in a single cell culture unit comprised of a mixture of cells grown in a manner to co-exist as a monolayer of relatively equivalent ratio and demonstrating complementary susceptibilities to a wider range of viruses and/or other organisms than could be detected by each individual cell line. For example, the viral assays involve
15 inoculating a cell mixture with a specimen suspected of containing a virus, allowing a sufficient period of time for the virus infectious cycle to proceed, followed by the detection and/or quantification of the number of virus-infected cells to determine the number of infectious virions in the specimen. This detection step may be accomplished using any number of available confirmation methods, including specific viral antigen
20 detection using antigen-specific antibodies, nucleic acid probes, and reporter gene detection. The assay also provides reliable methods and compositions for the quantification of the number of infectious virions present in a sample. In addition, the methods and compositions of the present invention are sufficiently sensitive that the presence of a single virion in a specimen may be detected.

25 The present invention also provides compositions comprising novel mixtures of various cell types traditionally used in single cell assays. In preferred embodiments, the cells are mixed to produce mixed monolayer cell cultures. One such mixed cell culture includes mink lung (*e.g.*, Mv1Lu) cells co-cultivated with human mucoepidermoid cells (*e.g.*, NCI-H292; also referred to as "H292" cells). This cell mixture is susceptible to
30 viruses such as influenza A, influenza B, RSV, parainfluenza types 1, 2, and 3, adenovirus, and CMV (*i.e.*, the group of viruses most commonly associated with respiratory virus disease). In other mixed cultures, buffalo green monkey kidney cells (BGMK) are co-cultivated with NCI-H292 cells for the detection and identification of enteroviruses, such as poliovirus, echoviruses and Coxsackie B virus (*e.g.*, Coxsackie A

and B viruses), and numbered EV strains. In addition to enteroviruses, it is contemplated that the present invention encompass cell types that are susceptible to picornaviruses such as Hepatitis A.

5 The present invention also provides compositions comprising novel mixtures of different cell types traditionally used in single cell assays that are co-cultivated with genetically engineered cells. In particularly preferred embodiments, the genetically engineered cell line is a DNA-transfected cell line that is susceptible to infection by a virus, the cell line having been stably transformed with a chimeric gene comprising a virus-inducible promoter and a gene coding for an enzyme, the expression of the enzyme
10 being dependent upon the presence of the virus. Such genetically engineered cells are, for example, described in U.S. Patent No. 5,418,132, herein incorporated by reference. In one preferred embodiment, a cell mixture includes human lung fibroblasts (*e.g.*, MRC-5 cells) co-cultivated with a stable baby hamster kidney (BHK) cell line, the genome of which has been engineered to contain the *E. coli lacZ* gene behind (*i.e.*, 3' to) an inducible
15 HSV promoter, HSV-1 ICP6 promoter (BHK/ICP6LacZ-5 cells are available from the ATCC as CRL-12072). This cell mixture is susceptible to infection by CMV and HSV types 1 and 2.

In yet another embodiment, the present invention provides compositions comprising novel mixtures of different types of genetically engineered cells. In
20 particularly preferred embodiments, the genetically engineered cell line is a DNA-transfected cell line that is susceptible to infection by a virus, the cell line having been stably transformed with a chimeric gene comprising a virus-inducible promoter and a gene coding for an enzyme, the expression of the enzyme being dependent upon the presence of the virus. The second genetically engineered cell line is a DNA-transfected
25 cell line susceptible to viral infection, which is stably transformed with a chimeric gene comprising a virus-inducible promoter, and a gene encoding a second enzyme (*i.e.*, an enzyme that is different from that associated with the first cell line), in which the expression of the second enzyme is dependent upon the presence of a second virus. In one preferred embodiment, a cell mixture is prepared in which engineered BHK cells
30 (*e.g.*, BHK/ICP6/LacZ-5 cells) are co-cultivated with a stable mink lung cell line (Mv1Lu), the genome of which has been engineered to contain the an inducible CMV promoter (the CMV UL45 promoter); these cells are referred to as "MLID5" cells, and are disclosed in U.S. Patent Appln. Ser. No. 08/846,026, herein incorporated by reference. This cell mixture is susceptible to infection by CMV and HSV virus types 1 and 2 (HSV-
35 1 and HSV-2), with CMV infecting the genetically engineered BHK cells, and HSV-1

and HSV-2 preferentially infecting the mink lung cells. In another embodiment, the present invention contemplates the use of genetically engineered cells (*e.g.*, mink lung cells) in which the cell genome is engineered to contain the firefly luciferase gene behind (*i.e.*, 3' to) an inducible CMV promoter; these cells are also described in U.S. Patent
5 Appln. Ser. No. 08/846,026. However, it is not intended that the present invention be limited to any particular cell types or cell lines, nor is it intended that the present invention be limited to any particular combinations of cells. It is also not intended that the present invention be limited in terms of the genetically engineered cells.

The following Table provides a matrix which indicates the ability of various cells
10 to form single, confluent monolayers, as well as co-cultivated, confluent, mixed cell monolayers.

Table 1. Cell Cultures

	MRC-5	CV-1	BGMK	McCoy	BHK*	A549	HEp-2	Mv1Lu	NCI-H292
	1	2	3	4	5	6	7	8	9
MRC-5	A	+	No	+	+	+	+	+	+
CV-1	B	++	No	+	+	+	+	+	+
BGMK	C		++	+	+	+	+	+	+
McCoy	D			++	+	+	Yes	+	+
BHK*	E				++	+	+	+	+
A549	F					++	+	+	+
HEp-2	G						++	+	+
Mv1Lu	H							++	+
NCI-H292	I								++

++ Denotes single cell types producing confluent monolayers
+ Denotes some degree of dimorphic, mixed monolayer
Yes Denotes cell mixtures that appear very uniform, with an even distribution
No Denotes cell mixtures that did not appear to work
* Denotes genetically engineered ELVIS BHK cells.

In yet another embodiment, the present invention provides kits for assaying samples for the presence of infectious viruses. In these kits, mixed cell cultures are provided which facilitate the detection and identification of particular virus groups (e.g., viruses associated with respiratory infections/diseases). In the kits, co-cultivated cells are supplied either frozen or dispensed (*i.e.*, ready for use) in shell vials, tubes, or multiwell plates. These cells are susceptible to infection by the virus group of interest as indicated by the sample type. In preferred embodiments, the kits also include reagents necessary to detect expression of viral antigens or virus-induced reporter gene expression.

One of the several advantages of the present invention is that it provides rapid and sensitive assay systems for the detection and identification of a single virus type from a multiplicity of possibilities, in a single mixed cell unit that is suitable for diagnostic assay. Thus, the present invention eliminates the need for multiple cell lines cultured in individual containers, provides reliable results in 1-3 days following inoculation of the cell cultures, rather than 1-28 days, eliminates the necessity of working with primary cell cultures, provides an efficient screening method for grouping and preliminary identification of viruses, and provides assay systems that are highly specific for viruses capable of inducing reporter gene expression. Thus, the present invention clearly fulfills a need that has been heretofore unmet in the field of diagnostic virology.

In a further embodiment, the invention provides a composition comprising a mixed cell culture comprising MDCK cells and one or more of A549 cells and H292 cells. These compositions are useful in detecting the presence of one or more of influenza viruses (such as influenza A and/or B), respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus. These methods are also useful in producing one or more of influenza viruses (such as influenza A and/or B), respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus.

The term "MDCK cells" and "Madin-Darby canine kidney cells" refer to cells that were isolated as previously described (Madin & Darby (1958) Tech. Prog. Rep. No. 25, Appendix VIII, p. 276. Naval Biological Laboratory, California and to cells that are established from these cells. MDCK cells are exemplified, but not limited to those deposited as ATCC accession number CCL-34. The term "established from" when made in reference to any cell disclosed herein (such as MDCK cell, A549 cell, H292 cell, *etc.*) refers to a cell which has been obtained (*e.g.*, isolated, purified, *etc.*) from the parent cell in issue using any manipulation, such as, without limitation, infection with virus, transfection with DNA sequences, treatment and/or mutagenesis using for example

chemicals, radiation, *etc.*, selection (such as by serial culture) of any cell that is contained in cultured parent cells. For example, the invention includes within its scope cell lines that may be established from any cell disclosed herein (such as MDCK cell, A549 cell, H292 cell, *etc.*) by treatment with chemical compounds (*e.g.*, *N*-ethyl-*N*-nitrosourea (ENU), methyl nitrosourea (MNU), procarbazine hydrochloride (PRC), triethylene melamine (TEM), acrylamide monomer (AA), chlorambucil (CHL), melphalan (MLP), cyclophosphamide (CPP), diethyl sulfate (DES), ethyl methane sulfonate (EMS), methyl methane sulfonate (MMS), 6-mercaptopurine (6MP), mitomycin-C (MMC), procarbazine (PRC), *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (MNNG), $^3\text{H}_2\text{O}$, and urethane (UR)), and electromagnetic radiation (*e.g.*, X-ray radiation, gamma-radiation, ultraviolet light).

Thus, reference to any virus or cell herein includes "wild-type" viruses and cells (*i.e.*, a virus or cell whose genome has not been manipulated by man) and "transgenic" viruses and cells (*i.e.*, a virus or cell that contains a heterologous nucleic acid sequence introduced into the virus or cell by means of molecular biological techniques).

Transgenic viruses and cells may contain heterologous nucleotide sequences such as reporter genes (such as *e.g.*, the *uid A* gene, β -glucuronidase gene, green fluorescent protein gene, *E. coli* β -galactosidase (LacZ) gene, *Halobacterium* β -galactosidase gene, *E. coli* luciferase gene, *Neurospora* tyrosinase gene, Aequorin (jellyfish bioluminescence) gene, human placental alkaline phosphatase gene, and chloramphenicol acetyltransferase (CAT) gene), transcriptional and translational regulatory sequences, selectable marker proteins (*e.g.*, proteins which confer drug resistance such as the bacterial aminoglycoside 3' phosphotransferase gene (also referred to as the neo gene) which confers resistance to the drug G418 in cells, the bacterial hygromycin G phosphotransferase (hyg) gene which confers resistance to the antibiotic hygromycin, and the bacterial xanthine-guanine phosphoribosyl transferase gene (also referred to as the gpt gene) which confers the ability to grow in the presence of mycophenolic acid, the HSV-tk gene and the dt gene), probe genes (such as the staphylococcal protein A and its derivative ZZ (which binds to human polyclonal IgG), histidine tails (which bind to Ni^{2+}), biotin (which binds to streptavidin), maltose-binding protein (MBP) (which binds to amylose), glutathione S-transferase (which binds to glutathione)), and receptor genes.

In one embodiment, equivalent cells within the scope of the invention include cells that are established from the exemplary MDCK cells deposited as ATCC accession CCL-34, and that have substantially the same sensitivity, increased sensitivity, or reduced

sensitivity to one or more of influenza virus A and influenza virus B as the cell from which it is established. The term "sensitivity" and "sensitive" when made in reference to a cell is a relative term which refers to the degree of permissiveness of the cell to a virus as compared to the degree of permissiveness of another cell to the same virus. For example, the term "increased sensitivity" to influenza virus when used in reference to the sensitivity of a first cell relative to a second cell refers to an increase in the first cell, preferably at least a 5%, more preferably from 5% to 10,000%, more preferably from 5% to 1,000%, yet more preferably from 10% to 200%, and even more preferably from 10% to 100%, increase in the quantity of influenza virus protein, influenza virus nucleic acid, and/or of CPE by progeny virus which is produced following infection of the first cell with influenza virus, as compared with the quantity of influenza virus protein, influenza virus nucleic acid, and/or of CPE by progeny virus (respectively) which is produced following infection of the second cell. For example, if 34 samples containing influenza virus were tested for the presence of progeny virus, with 25 and 13 samples showing the presence of CPE using a first cell and second cell, respectively, then the sensitivity is 74% and 38% for the first cell and second cell, respectively. This reflects an increase of 90% in the sensitivity of the first cell as compared to the sensitivity of the second cell.

In another embodiment, equivalent cells within the scope of the invention include cells that are established from the exemplary MDCK deposited as ATCC accession number CCL-34, and that have substantially the same sensitivity to influenza virus as the cell from which it is established. This may be advantageous where, for example, the parent cell is made transgenic for a reporter gene.

In a further embodiment, equivalent cells within the scope of the invention include cells that are established from the exemplary MDCK deposited as ATCC accession number CCL-34, and that have increased sensitivity or decreased sensitivity to influenza virus as compared to cells from which they were established. This may be desirable where, for example, the parent cell is made transgenic for a receptor gene which alters the level of binding of influenza B virus to the cell.

The invention's methods that employ mixed cell cultures containing MDCK cells are useful for detecting influenza virus. The term "detecting" when in reference to detecting the presence of any virus in cells refers to determining the presence, using any method, of the virus inside the cells, on the cells, and/or in the medium with which the cells come into contact. These methods are exemplified by, but not limited to, the observation of cytopathic effect, detection of viral protein, such as by

immunofluorescence and Northern blots, and detection of viral nucleic acid sequences, such as by PCR, reverse transcriptase PCR (RT-PCR), Southern blots and Northern blots.

As used herein the term "influenza virus" refers to members of the orthomyxoviridae family of enveloped viruses with a segmented antisense RNA genome (Knipe and Howley (eds.) Fields Virology, 4th edition, Lippincott Williams and Wilkins, Philadelphia, PA [2001]). Two types of influenza virus (A and B) are human pathogens causing respiratory pathology.

When A549 and/or H292 cells are in mixed cell culture with MDCK cells, the mixed cell cultures may also be used to detect and propagate other viruses than influenza virus, such as respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus.

The terms "respiratory syncytial virus" and "RSV" refer to one or more members of the family Paramyxoviridae, subfamily pneumovirus, which are enveloped, single stranded antisense RNA viruses that infect the respiratory tract (Schmidt and Emmons(eds.) Diagnostic Procedures for Viral, Rickettsial and Chlamydial Infections, 6th edition, American Public Health Assoc. Inc.[1989] Washington, DC. There are two major strains of RSV represented by, but not limited to, Long (Group 1)ATCC VR-26, and 18537 (Group 2)ATCC VR-1401. The following five exemplary human RSV strains are available from ATCC: VR-1400, VR-1401, VR-1540, VR-26, and VR-955.

As used herein, the term "parainfluenza virus" refers to certain members of the paramyxoviridae family of enveloped viruses with a single-stranded antisense RNA genome (Knipe and Howley (eds.) Fields Virology, 4th edition, Lippincott Williams and Wilkins, Philadelphia, PA [2001]). Four types of parainfluenza virus (1 to 4) are human respiratory pathogens. Prototype strains of the human paramyxoviruses parainfluenza types 1, 2, 3, 4A, 4B, and mumps, may be obtained from the reference virus collection of the Respiratory and Enteric Viruses Branch of the Center for Infectious Diseases, Centers for Disease Control ("CDC"), Atlanta, Ga (see U.S. Patent 5,262,359 to Hierholzer). These strains are also available from the ATCC, Rockville, Md., under accession numbers VR-94, VR-92, VR-93, VR-279, VR-579, and VR-106, respectively (see U.S. Patent 5,262,359 to Hierholzer).

As used herein, the term "adenovirus" refers to a double-stranded DNA adenovirus of animal origin, such as avian, bovine, ovine, murine, porcine, canine, simian, and human origin. Avian adenoviruses are exemplified by serotypes 1 to 10 which are available from the ATCC, such as, for example, the Phelps (ATCC VR-432), Fontes (ATCC VR-280), P7-A (ATCC VR-827), IBH-2A (ATCC VR-828), J2-A (ATCC

VR-829), T8-A (ATCC VR-830), and K-11 (ATCC VR-921) strains, or else the strains designated as ATCC VR-831 to 835. Bovine adenoviruses are illustrated by those available from the ATCC (types 1 to 8) under reference numbers ATCC VR-313, 314, 639-642, 768 and 769. Ovine adenoviruses include the type 5 (ATCC VR-1343) or type 6 (ATCC VR-1340). Murine adenoviruses are exemplified by FL (ATCC VR-550) and E20308 (ATCC VR-528). Porcine adenovirus (5359) may also be used. Adenoviruses of canine origin include all the strains of the CAV1 and CAV2 adenoviruses [for example, Manhattan strain or A26/61 (ATCC VR-800) strain]. Simian adenoviruses are also contemplated, and they include the adenoviruses with the ATCC reference numbers VR-591-594, 941-943, and 195-203. Human adenoviruses, of which there greater than fifty (50) serotypes are known in the art, are also contemplated, including the Ad2, Ad3, Ad4, Ad5, Ad7, Ad9, Ad12, Ad17, and Ad40 adenoviruses.

The terms "metapneumovirus" and "MPV" refer to a a negative-sense single stranded RNA virus belonging to the Paramyxoviridae family, subfamily Pneumovirinae, and genus Metapneumovirus. MPV includes mammalian MPV, which is exemplified by human, primate, horse, cow, sheep, pig, goat, dog, cat, avian and rodents MPV. Mammalian MPV is phylogenetically more closely related to particular virus isolates than to turkey rhinotracheitis virus, the etiological agent of avian rhinotracheitis, and also may be identified by its genomic organization (see, for example, U.S. patent application publication numbers 20030232326, 20040005544, 20040005545, and 20030232061, and published WO 02057302A2 and WO 03072719A2). The invention contemplates each of the variant MPV that are idnetified based on the relative homology of their genomic sequences to other viruses, as described in, for example, U.S. patent application publication numbers 20030232326, 20040005544, 20040005545, and 20030232061, and published WO 02057302A2 and WO 03072719A2.

MPV may be detected by, for example, detecting cytopathic effect in the exemplary LLC-MK2 cells and HEp-2 cells (Chan et al. 2003 Emerging Infectious Diseases, 9:1058-1063; Setterquist et al., 19th Annual Clinical Virology Symposium, Clearwater Florida, April 27-30, 2003), detecting MPV proteins using antibodies and/or detecting MPV nucleic acid sequences (see, for example, U.S. patent application publication numbers 20030232326 and 20040005544). In one embodiment, MPV nucleic acid sequences may be detected in the absence of detectable CPE.

The invention's data is the first demonstration of the use of MDCK in mixed cell culture (Examples 4-9), and is contrasted with Frank et al. (1979) Journal of Clinical Microbiology, 10(1):32-36 which disclosed using MDCK cells. The ability to grow

MDCK in mixed cell culture with the exemplary cell lines of H292 and A549 was surprising in view of data herein (Example 1) which demonstrates the unpredictability of co-culturing two or more cell lines, as well as the unpredictability that once co-cultured, the cells will retain their biological activity in detecting and/or producing virus.

One advantage of using MDCK cells in the invention's mixed cell cultures with A549 and/or H292 is that these cells are non-permissive to SARS-CoV infection as determined by CPE (Table 12 herein; see also Drosten, et al., 2003, N.Engl.J.Med. 348:1967-1976; Ksiazek, et al., 2003, N.Engl.J.Med. 348:1953-1966; Peiris, et al., 2003, Lancet 361:1319-1325). Thus, an advantage of using MDCK cells is that they permit detection of respiratory viruses (such as respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, parainfluenza 3 virus, and metapneumovirus), while being nonpermissive, or having a low level of permissivity, to SARS-CoV (Table 12). Thus, mixed cell cultures containing MDCK are useful for increasing the safety of cell cultures that are used in screening clinical samples for respiratory pathogens other than SARS-coronavirus. This is particularly useful in small laboratories that detect respiratory viruses (such as respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus, and parainfluenza 3 virus), because the use of mixed cell cultures containing MDCK by these laboratories would obviate the need to resort to containment approaches that would otherwise be required for cells producing infectious SARS-CoV.

In particular, although both MDCK and Mv1Lu cells are susceptible to influenza B virus (Example 2), data herein shows, surprisingly, that MDCK has a substantially lower level of permissivity and/or susceptibility to SARS-CoV as compared to Mv1Lu (Table 12). The terms "lower," "smaller," "reduced," "decreased," and grammatical equivalents when in reference to the level of permissivity and/or susceptibility to a virus by a first cell type relative to a second cell type, mean that the level of permissivity and/or susceptibility to the virus in the first cell type is lower than in the second cell type by any amount that is statistically significant, using any art-accepted statistical method of analysis. In one embodiment, the level of permissivity and/or susceptibility to the virus in the first sample is at least 10% lower than, at least 25% lower than, at least 50% lower than, at least 75% lower than, at least 85% lower than, at least 90% lower than, at least 95% lower than, and/or at least 99% lower than the level of permissivity and/or susceptibility to the virus in the second cell type. Data herein shows that, in one embodiment, the level of permissivity and/or susceptibility of MDCK cells to SARS-CoV is 0.004% the level of susceptibility of Mv1Lu cells (Table 12).

The terms "SARS coronavirus," "SARS-CoV," and "severe acute respiratory syndrome coronavirus" are equivalent, and are used to refer to an RNA virus that is the causative agent of severe acute respiratory syndrome (Drosten, et al., 2003, *supra*; Fouchier, et al., 2003, *supra*; Ksiazek, et al., 2003, *supra*; Peiris, et al., 2003, *supra*; Poutanen, et al., 2003, *supra*). Exemplary strains of SARS coronavirus include, but are not limited to, Urbani, Tor2, CUHK-W1, Shanhgai LY, Shanghai QXC, ZJ-HZ01, TW1, HSR 1, WHU, TWY, TWS, TWK, TWJ, TWH, HKU-39849, FRA, TWC3, TWC2, TWC, ZMY 1, BJ03, ZJ01, CUHK-Su10, GZ50, SZ16, SZ3, CUHK-W1, BJ04, AS, Sin2774, GD01, Sin2500, Sin2677, Sin2679, Sin2748, ZJ-HZ01, and BJ01.

However, coronaviruses can establish persistent infection in cells without inducing CPE, suggesting that CPE may not be an accurate indicator of infection (Chaloner, et al., 1981, Arch.Virol. 69:117-129). Data herein confirmed this surprising observation by demonstrating replication of SARS-CoV in the absence of CPE. For example, Example 12 shows replication of SARS-CoV, as detected by sgRNA and virus titers, in the absence of CPE. In particular, significant CPE was not observed in pRhMK, pCMK, R-mix (Mv1Lu and A549), Mv1Lu, HEK-293T, and Huh-7 cells at 5 days post infection, although virus titers as well as SARS-CoV sgRNA were actually increased within 24 hours post infection (Table 12).

The terms "subgenomic RNA" and "sgRNA" are used interchangeably herein to refer to a nucleotide sequence comprising at least a portion of the leader sequence of SARS-CoV.

The term "leader sequence" refers to a sequence of about 40 to about 150, about 50 to about 80, and or about 55 to about 75, nucleotides that is located at the 5' terminus of the genome. This sequence is juxtaposed to the 5' terminus of each subgenomic RNA by transcriptional mechanisms during synthesis. There is very strong sequence conservation of the leader sequence across the strains of SARS. In one embodiment, the leader sequence is exemplified by the sequence from nucleotide 1 to nucleotide 72 for SARS-CoV (Urbani) 5'-atattaggtttttacctaccaggaaaagccaaccaacctcgatctc ttgtagatctgttctctaaacgaac-3' (SEQ ID NO:1); 5'-tattaggtttttacctaccaggaaaag ccaaccaacctcgatctctttagatctgttctctaaacgaac-3' (SEQ ID NO:2) of gi|33304219|gb|AY351680.1| SARS coronavirus ZMY 1, 5'- taggtttttacctaccaggaaaagccaaccaacctcgatctctttagatctgttctctaaacgaac-3' (SEQ ID NO:3) of gi|31416305|gb|AY278490.3| SARS coronavirus BJ03, 5'- ctaccaggaaaagccaaccaacctcgatctctttagatctgttctctaaacgaac-3' (SEQ ID NO:4) of gi|30421451|gb|AY282752.1| SARS coronavirus CUHK-Su10, 5'-

taccaggaaaagccaaccaacctcgatctctttagatctgttctctaaacgaac-3' (SEQ ID NO:5) of
gi|31416306|gb|AY279354.2| SARS coronavirus BJ04, and 5'-
ccaggaaaagccaaccaacctcgatctctttagatctgttctctaaacgaac-3' (SEQ ID NO:6) of
gi|30275666|gb|AY278488.2| SARS coronavirus BJ01.

5

EXPERIMENTAL

The following examples are provided in order to demonstrate and further illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

10

In the experimental disclosure which follows, the following abbreviations apply:

eq (equivalents); M (Molar); μ M (micromolar); N (Normal); mol (moles); mmol (millimoles); μ mol (micromoles); nmol (nanomoles); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); l or L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); xg (times gravity); $^{\circ}$ C (degrees Centigrade); FBS (fetal bovine serum); PBS (phosphate buffered saline; HEPES (N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid]); HBSS (Hank's Balanced Salt Solution); MEM (Minimal Essential Medium); EMEM (Eagle's Minimal Essential Medium); BBL (Becton Dickinson Microbiology Systems, Cockeysville, MD); DIFCO (Difco Laboratories, Detroit, MI); U.S. Biochemical (U.S. Biochemical Corp., Cleveland, OH); Chemicon (Chemicon, Inc., Temecula, CA); Dako (Dako Corporation, Carpinteria, CA); Fisher (Fisher Scientific, Pittsburgh, PA); Sigma (Sigma Chemical Co., St. Louis, MO.); ATCC (American Type Culture Collection, Rockville, Maryland); Bartel's (Bartels, Issaquah, WA); and BioWhittaker (BioWhittaker, Walkersville, MD).

15

20

25

The cells used during the development of the present invention and described in the following Examples, were obtained from the ATCC, with the exception being that the BGMK and PRMK cells were obtained from BioWhittaker, and the MRC-5 cells were obtained from both ATCC and BioWhittaker. The ATCC numbers of the cells are indicated in the following Table.

Table 2. ATCC Cell Lines

	Cell Line	ATCC Number
	BHK/ICP6LacZ-5	CCL-12072
	A549	CCL-185
5	CV-1	CCL-70
	HEp-2	CCL-23
	hs27	HFF; CRL-1634
	Mv1Lu	CCL-64
	McCoy	CCL-1696
10	NCI-H292	CCL-1848
	MRC-5	CCL-171
	WI-38	CCL-75
	Vero	CCL-81
	MDCK (NBL-2)	CCL-34
15	BHK21	CCL-10
	HEL299	CCL-137
	HeLa	CCL-2
	Mv1Lu-hF	PTA-4737

EXAMPLE 1

Co-Cultivation of Cells

In this Example, mixed cell cultures were established in which single, dimorphic cell sheets were produced at confluency.

5 In these experiments, all of the cell lines were cultured to confluency in sterile polystyrene flasks in EMEM (Eagle's Minimal Essential Medium) with 25 mM HEPES, 7% fetal bovine serum (FBS), 2 mM L-glutamine, and penicillin/streptomycin (100 Units/100 µg per ml of medium each).

10 Cells to be cultured were harvested by first rinsing source cell monolayers with Hank's Balanced Salt Solution (HBSS) without magnesium or calcium. Depending upon the cell line, the cells were dissociated by adding trypsin (0.125% in HBSS, without calcium or magnesium) or trypsin-EDTA (0.25% in 1 mM EDTA in HBSS, without calcium or magnesium), or directly to the cell monolayer, and incubating for approximately 5 minutes at ambient temperature. Ten volumes of cell culture medium
15 was added to each trypsinized cell suspension and the cells were repeatedly pipetted in order to produce near-single cell suspensions (*i.e.*, without cell aggregates). Each trypsinized cell suspension was diluted in an adequate volume of culture medium to produce an optical density of cell suspension suitable to produce a confluent monolayer of cells within 2-3 days of incubation in a 96-well microtiter plate. For single cell
20 monolayers (*i.e.*, one cell type per well), 0.2 ml of suspension was used to inoculate each well. For example, the final cell preparations ranged from a final optical density at 500 nm of 0.012 OD units/ml for CV-1 cells to 0.03 OD units/ml for HEp-2 cells.

Cell mixture monolayers were produced by co-planting two distinct cell types at an equal volume of each diluted cell suspension (*i.e.*, 0.1 ml of each cell type was used to
25 inoculate each well of a 96-well microtiter plate). The cells were allowed to attach to the well surface by gravity for 30-60 minutes, and the inoculated microtiter plates were incubated for up to three days at 36°C in 5% CO₂ with 95% relative humidity.

Periodically during incubation, single and mixed monolayers were checked for overall viability. The mixed cell culture monolayers were also checked for the ability of
30 the cell lines to co-exist and develop as a single cell sheet (*i.e.*, a single monolayer), with two distinct cell morphologies (*i.e.*, dimorphic cell sheets), at an approximately equal density of each cell type. At confluency, the cells were treated with a methylene blue staining solution to fix the cells and stain them a light blue in order to provide contrast for visualization using light microscopy.

Some of the mixed monolayers successfully grew as a mixed cell monolayer adhered to the well surfaces, exhibiting a smooth, evenly distributed monolayer. These mixed cultures were designated as "morphologic category 1." In these cultures, each cell type could be easily distinguished and appeared to survive well in a mixed monolayer, giving the appearance of a single cell distribution. Mixed monolayers composed of HEp-2 and McCoy cells displayed this morphology.

Some of the mixed monolayers successfully grew as a mixed monolayer adhered to the well surfaces, but exhibited two distinct morphologies at confluency. These mixed cultures were designated as "morphologic category 2." In these cultures, separate, distinct patches of each cell line co-existed within the monolayer, giving the appearance of oil mixing with water. Although an understanding of the mechanism is not necessary in order to use the present invention, it is likely that this appearance is most likely the result of contact inhibition between two specific cell types. The relative sizes of the patches was found primarily to be a function of how evenly the cells were distributed at cell planting. The more even the cell distribution at planting, the patches or islands were smaller as the monolayer reached confluency. Examples of monolayers that produced this appearance were mink lung cells co-cultivated with NCI-H292 cells, mink lung cells co-cultivated by buffalo green monkey kidney (BGMK) cells, and human lung carcinoma A549 cells co-cultivated with NCI-H292 cells.

However, some cells types could not produce a mixed cell monolayer, when mixed at relatively equal cell numbers at planting in the same culture medium. In some of these cultures, only one of the cell types was found to be viable (*i.e.*, the culture was effectively a single cell type). Examples of mixed cell cultures that were found to be unsuitable for the production of mixed monolayers include human embryonic lung fibroblasts (MRC-5 cells) co-cultivated with BGMK cells. In this mixture, the MRC-5 cells become toxic and form aggregates of dead cells soon after planting. Thus, at confluency, the monolayer only contains one functional, viable cell type, the BGMK cells. Thus, this cell mixture was found to be unsuitable for producing mixed cell monolayers as the cells failed to form mixed cell monolayers of either a smooth or dimorphic morphologic type.

EXAMPLE 2

Detection of Respiratory Viruses in Mixed Cell Cultures

5 In this Example, mixed cell cultures were used to detect various respiratory viruses including Influenza A, RSV, adenoviruses, parainfluenza viruses, and Influenza B, present in clinical specimens. The mixed cells used in these experiments were Mv1Lu (mink lung cells) and NCI-H292 (human mucoepidermoid cells).

Cell Lines

10 Confluent T-225 flasks of Mv1Lu and H292 cells were prepared in EMEM with HEPES, 10% FBS, 2 mM L-glutamine, and 50 µg/ml gentamicin. The cells were harvested by first rinsing them in 30 ml HBSS without magnesium and calcium. The cells were then dissociated from the flask by brief exposure (*i.e.*, until the cells lifted from the bottom of the flask) to 7 ml trypsin-EDTA solution as described in Example 1. Then, 30 ml media was added to the cells to prepare a cell suspension concentrate. The
15 optical density of each cell suspension was determined at 500 nm, using 3 ml of cells. Typically, the OD reading was 0.2/ml for both the Mv1Lu and H292 cells. In addition to the Mv1Lu and H292 cells, rhesus monkey kidney cells (PRMK), A549 cells, and MDCK cells were used in the present Example. These additional cell lines were prepared in single cell cultures as known in the art.

20 Mixed Cell Cultures

When each cell suspension concentrate was determined to be 0.2 OD units/ml, 5.2 ml of the Mv1Lu, and 8.7 ml H292 cell suspensions were added to 86.1 ml of culture medium, in order to provide an acceptable working ratio of each cell type (*i.e.*, it was a preparation of diluted mixed cells). This ratio was devised in order to achieve a confluent
25 monolayer, in which each cell type covered a substantially equivalent surface area within 1-3 days post-planting of the diluted mixed cells. Prior to dispensing, care was taken to prepare homogenous suspensions of diluted mixed cells. The mixed cells were dispensed at 0.75 ml per glass shell vial (*i.e.*, glass vial containing a sterile glass coverslip). After
30 planting, the vials were allowed to sit for 60 minutes at ambient temperature so that the cells could settle by gravity and produce a more optimum cell distribution of each cell type. The mixed cells were then incubated for 1-3 days at 36°C in 5% CO₂, at 95% relative humidity. Subsequently, the shell vials were stored at ambient temperature to

maintain each cell type at substantively equivalent surface ratios for up to 10 days from achieving confluency.

Samples and Processing

Nasopharyngeal specimens submitted to a diagnostic virology laboratory were obtained from patients exhibiting influenza-like symptoms. The specimens were centrifuged to produce a cell pellet for direct antigen testing, and a specimen supernatant for inoculation of various cell cultures. The cell pellet was resuspended in phosphate buffer to prepare a cell suspension and 25 µl portions of the cell suspension were spotted onto a glass slide and dried. Each spot of cells on the slide were then fixed with fixative (e.g., acetone), and incubated for 30 minutes with individual antibody solutions (Bartel's) capable of recognizing various respiratory viruses, including influenza A and RSV, as well as other respiratory viruses. A second antibody solution containing fluorescein (FITC) labelled goat anti-mouse antibodies and counterstain (Bartel's) was added to cover each cell spot on the slides, and incubated for an additional 30 minutes at 35-37°C. The counterstain in the FITC-goat anti-mouse antibody solution contains Evans Blue, which stains the cells and appears red under fluorescence. Slides prepared from the nasopharyngeal specimens were observed for positive (*i.e.*, virus-infected), apple green staining fluorescent cells, using epifluorescence at 100-400X magnification.

In addition, 0.2 ml aliquots of the specimen supernatant were inoculated onto various cell cultures prepared in shell vials containing glass coverslips. The cell cultures included primary rhesus monkey kidney cells (PRMK; ViroMed or BioWhittaker), Mv1Lu cells (Diagnostic Hybrids) HEp-2 cells (Diagnostic Hybrids), MDCK, A549, and H292 cells, as single cell monolayers, as well as mixed cell monolayers of Mv1Lu and H292 cells, produced as described above.

Each inoculated shell vial was centrifuged for 60 minutes at 700 xg, and then incubated for 1-3 days at 36°C, in appropriate culture medium (e.g., EMEM containing 0.5 to 2% FBS, 2 mM L-glutamine, and penicillin/streptomycin [100 Units/100 µg per ml of medium each]). After incubation, the culture medium was decanted, and the cells were fixed to the glass coverslip with a solution of acetone and methanol (50:50, v/v). An antibody solution (Chemicon or Bartel's) containing a pool of monoclonal antibodies to multiple respiratory viruses, including Influenza A and RSV, as well as other respiratory viruses was added to cover each coverslip. The coverslips were then incubated for 30 minutes at 35-37°C. The antibody solution was then removed and the coverslips were rinsed with PBS. A second antibody solution containing fluorescein (FITC) labelled goat

anti-mouse antibodies and counterstain (Bartel's) was added to cover each coverslip, and incubated for an additional 30 minutes at 35-37°C. The counterstain in the FITC-goat anti-mouse antibody solution contains Evans Blue, which stains the cells and appears red under fluorescence. Shell vial coverslips prepared from the nasopharyngeal specimens (*i.e.*, inoculated cultures) were observed for positive (*i.e.*, virus-infected), apple green staining, fluorescent cells, using epifluorescence at 100-400X magnification.

Results

Some specimens demonstrated a positive direct antigen reaction on the cell spot incubated with Influenza A monoclonal antibody. These specimens also demonstrated fluorescent staining on the single cell Mv1Lu coverslip and the Mv1Lu/H292 mixed cell coverslip, but no or very little fluorescence on the single cell H292 coverslip. The H292 cells are either not susceptible to this strain of Influenza A, or are significantly less susceptible, such that infection is not detectable. Additionally, in some cases (*i.e.*, in specimens with low virus titers), the culture systems were more sensitive than the direct antigen detection method. Also, while the single PRMK cell cultures (*i.e.*, the "gold standard" cells used to detect Influenza A) were positive for the presence of Influenza A, with many specimens, the numbers of infected cells and the total of number of positive specimens were lower than those identified as positive by the mixed cell monolayers.

In addition, both the MDCK and PRMK cells missed one low titer specimen positive for Influenza A by direct antigen testing (IFA), and one other specimen that was also positive for Influenza A by IFA, while the Mv1Lu cells detected the virus in all of the samples determined to be positive based on direct antigen detection (IFA).

Some specimens demonstrated a positive direct antigen reaction on the cell spot incubated with RSV monoclonal antibody. These specimens also demonstrated fluorescent staining on the single cell H292 coverslip and the MV1Lu/H292 mixed cell coverslip, but no or very little fluorescence on the single cell MV1Lu coverslip. H292 cells are susceptible to RSV infection, while Mv1Lu cells are not susceptible (or are significantly less susceptible, such that infection is not detectable). In addition to the mixed cell cultures, HEp-2 cells (*i.e.*, the "gold standard" cells used to detect RSV) were also observed for the presence of RSV; the performance of HEp-2 cells was generally less sensitive than that of the Mv1Lu and H292 mixed cell monolayers, or the H292 single cell monolayers. These results with Influenza A tested in mink lung cells was very surprising, as the detection of Influenza A using these cells has previously not been described.

Adenoviruses identified from five clinical specimens based on direct antigen testing (IFA) were detected in the H292 and cell culture mixes, while the PRMK cells missed two of the low titer specimens (*i.e.*, there were two false negatives). Thus, H292 and the mixed cultures were more sensitive than PRMK for detection of adenoviruses.

5 While the A549 cells may provide slightly more positive cells, the 292 cells, mixed cell cultures, and A549 cells detected an equal number of positive specimens.

Parainfluenza viruses were also detected in the H292 and mixed cell cultures, while the PRMK cells missed one low titer specimen.

10 These results clearly show that the mixed cell cultures were equal in sensitivity to the single cell (H292 and Mv1Lu) cultures. Thus, the mixed cells provide savings in material, time, space, and labor, while providing the same level of sensitivity in the detection of respiratory viruses as single cell cultures presently commonly used in diagnostic virology laboratories.

Influenza B Specimens

15 In addition to the samples discussed above, various dilutions of multiple Influenza B strains obtained from the ATCC were tested in MDCK, Mv1Lu, and PRMK cells. The following Table provides the results of these experiments. In this Table, "MD" refers to the "Maryland" strain, "HK" refers to the "Hong Kong" strain, "TW" refers to the "Taiwan" strain, and "MA" refers to the "Massachusetts" strain.

**Table 3. Comparison of Influenza B Virus Detection From
Prototype Viruses by MDCK, ML, and PRMK Cells**

Influenza B Virus Strain	Virus Dilution	Cell Line		
		MDCK	Mv1Lu	PRMK
MD	10 ⁻⁴	+	+	+
	10 ⁻⁵	+	+	+
	10 ⁻⁶	-	+	-
HK	10 ⁻⁴	+	+	+
	10 ⁻⁵	+	+	-
	10 ⁻⁶	-	-	-
TW	10 ⁻⁴	+	+	+
	10 ⁻⁵	+	+	+
	10 ⁻⁶	-	-	-
MA	10 ⁻⁴	+	+	+
	10 ⁻⁵	+	+	+
	10 ⁻⁶	+	+	+

These results indicate that Mv1Lu, MDCK, and PRMK are comparable for the detection of multiple Influenza B virus strains. Thus, these cell lines were identified as good candidates for mixed cell cultures, as well as single cell cultures for the identification of this virus.

EXAMPLE 3

Detection of CMV in Mixed Cell Cultures

In this Example, mixed cell cultures of Mv1Lu and NCI-H292 cells were used to detect the presence of human cytomegalovirus (HCMV).

The Towne strain of HCMV (ATCC #VR977) was amplified in MRC-5 cells to a titer of greater than 10⁶/ml, and frozen at -85°C in EMEM containing 10% FBS. Serial dilutions of HCMV were prepared and inoculated into single monolayers of mink lung (Mv1Lu) cells, MRC-5 cells, and mixed cell monolayers of Mv1Lu and H292 cells. Each infected cell culture system was centrifuged for 60 minutes at 700 xg, and then incubated for 24 hours at 36°C in 5% CO₂, in appropriate culture medium (e.g., EMEM containing

10% FBS). The culture medium was removed and the cells were fixed to the glass coverslip using a solution of 80% acetone in water. A sufficient amount of HCMV antibody solution (Chemicon) was added to cover each coverslip and incubated for 30 minutes at 35-37°C. The antibody solution was removed, and the coverslip was rinsed with PBS. A second antibody solution consisting of FITC-labelled goat anti-mouse antiserum was added to cover each coverslip and incubated an additional 30 minutes at 35-37°C. The specimens were then observed under epifluorescence at 100-400X magnification for positive (*i.e.*, CMV-infected), nuclear staining, fluorescent cells.

As described in previous Examples, the counterstain in the FITC-labelled goat anti-mouse antibody solution contains Evans Blue, which stains the cells and appears red, when excited by fluorescent light. Fluorescent, apple green nuclear stain was observed in the Mv1Lu single cell monolayer and in the mixed cell monolayers, but not in the H292 cells, as the Mv1Lu cells are susceptible to HCMV infection, while H292 cells are not (or the H292 cells are significantly less sensitive). The MRC-5 cells (*i.e.*, the "gold standard" cells for detection of HCMV) performed about the same as the mixed cell monolayer, as these cultures had a similar number of infected cells as the cells in the mixed monolayer.

EXAMPLE 4

Detection of Enteroviruses in Mixed Cell Cultures

In this Example, mixed cell cultures were used to detect the enteroviruses, Coxsackie B virus and Echovirus. In these experiments, a mixed cell monolayer of BGMK and NCI-H292 cells were used.

Confluent T-225 flasks of BGMK and H292 cells were prepared in EMEM with 25 mM HEPES, 10% FBS, 2 mM L-glutamine, and 50 µg/ml gentamicin. The cells were harvested by first rinsing in 30 ml HBSS without magnesium and calcium, and were then dissociated from the flasks by a brief treatment of 7 ml trypsin-EDTA solution (as described in Example 1). Then, an additional 30 ml of culture medium (EMEM with HEPES, 10% FBS, 2 mM L-glutamine, and 50 µg/ml gentamicin) was added to the suspension to produce a cell suspension concentrate. The optical density at 500 nm was determined for each suspension, using 3 ml of cells. Typically, the OD reading was 0.2/ml for both the BGMK and H292 cell suspensions.

Next, 3 ml of BGMK cell suspension and 8 ml of H292 cell suspension (both suspensions were at 0.2 OD units/ml) were then added to 29 ml of the culture medium (25 mM HEPES, 10% FBS, 2 mM L-glutamine, and 50 µg/ml gentamicin) to provide an acceptable working ratio of each cell type in a diluted mixed cell suspension. This ratio

was intended to achieve a confluent monolayer consisting of each cell type covering substantially equivalent surface area within 1-3 days post-planting of the diluted mixed cells. Care was exercised to prepare a homogenous suspension of diluted mixed cells prior to dispensing 0.75 ml to each of 100 glass shell vials, each of which contained a sterile glass coverslip. The vials were allowed to sit for 60 minutes post-planting at ambient temperature to allow the cells to settle by gravity and produce a more optimum cell distribution. The vials were then moved to an incubator for incubation at 36°C for 1-3 days in 5% CO₂, at 95% relative humidity.

Stock virus suspensions and clinical specimens shown to contain Coxsackie B virus or echovirus were used to infect BGMK/H292 cell mixtures, as well as single cell monolayers of BGMK, H292, MRC-5, and PRMK cells. For clinical samples, throat swab, nasopharyngeal swab, sputum, stool, and rectal swabs were collected from patients, placed in viral transport medium, and filtered through 0.45 µm filter to remove possible bacterial and fungal contaminants prior to inoculation of cell cultures. Cerebrospinal fluid (CSF) collected from patients was placed in viral transport medium, and used directly for inoculation of cells. For inoculation of shell vials, the media present in the vials were removed and fresh media added. Then, 0.2 ml of specimen was inoculated into each vial. The inoculated vials were centrifuged at 700 xg for 45-60 minutes at room temperature. Subsequently, the vials were incubated at 37°C for 1-3 days, and viral presence was detected using immunofluorescent staining.

For staining, the medium was removed from each vial and the cells were fixed on the coverslip with acetone. The coverslip was removed from each vial, and stained with 25 µl primary antibody (mouse monoclonal antibody directed against enteroviruses [Dako]), for 30 minutes at 37°C. After washing with PBS, 25 µl of the FITC-conjugated anti-mouse Ig (Dako) was used as a secondary antibody for staining, and incubated at 37°C for 30 minutes. After another wash, the coverslips were mounted on slides and observed under fluorescence. The presence of one or more specific fluorescent-stained cells on the coverslip was considered to be a positive. As described in previous Examples, the counterstain in the FITC-labelled goat anti-mouse antibody solution contains Evans Blue, which stains the cells, and appears red upon exposure to fluorescent light. For Coxsackie B virus detection, fluorescent, apple green stain was observed in many of the BGMK cells in the BGMK single cell monolayer and in the mixed cell monolayers primarily in the BGMK cells, but not in as many H292 cells, as BGMK cells are more susceptible to Coxsackie B virus infection. For some types of Coxsackie B virus isolates, H292 cells are not as susceptible (or the H292 cells are significantly less

susceptible). The "gold standard" cell line (*i.e.*, PRMK cells) did not exhibit the same number of infected cells as the mixed cell monolayers.

For detection of echovirus, fluorescent, apple green stain was observed in many H292 cells in the H292 single cell monolayer and in the mixed cell monolayers, primarily in the H292 cells, but not in as many BGMK cells, as H292 cells are more susceptible to echovirus infection, while BGMK cells are not as susceptible (or the BGMK cells are significantly less sensitive). The "gold standard" line (*i.e.*, MRC-5 cells) performed, but did not appear to have as many infected cells as the mixed cell monolayers. In the case of the BGMK/H292 mixed cell monolayers infected with high titer samples of enteroviruses, cell-specific virus mediated cytopathic effect (CPE) was evident (*i.e.*, the CPE was observed in BGMK cells when Coxsackie B virus was present at high titer, and CPE was observed in H292 cells when echovirus was present at high titer).

EXAMPLE 5

Detection of Herpes Simplex Virus and HCMV in Mixed Cell Cultures

In this Example, mixed cell cultures are used to detect herpes simplex virus (HSV) and HCMV, using a mixed cell monolayer of genetically engineered baby hamster kidney (BHK) cells (*e.g.*, ATCC #CCL-12072) and Mv1Lu cells.

The BHK and Mv1Lu cells are grown in flasks, trypsinized, and mixed as described in previous Examples, such that a suitable dilution of mixed cells is produced. These mixed cell dilutions are then used to inoculate sterile glass shell vials containing coverslips, as described above. The cells are then centrifuged and inoculated with virus or clinical samples, incubated, and fixed, as described above.

HCMV is detected in the Mv1Lu cells, using antibody as described in Example 3 above, and HSV (HSV-1 and HSV-2) are identified using a β -galactosidase staining kit (*i.e.*, detecting the reporter gene induced by the virus infecting the genetically engineered BHK cells).

EXAMPLE 6

Detection of Respiratory Viruses in Mixed Cell Cultures

In this Example, mixed cell cultures are used to detect a panel of respiratory viruses. In these experiments, three cell types are combined so as to produce a mixed cell culture that is capable of detecting at least three viruses.

First, A549, H292, and mink lung (*e.g.*, Mv1Lu) cells are grown in flasks, trypsinized, and mixed as described in previous Examples, such that a suitable dilution of mixed cells is produced. In preferred embodiments, the cells are diluted such that the mixed cells in culture will be in approximately the same proportions (*i.e.*, 1:1:1). These mixed cell dilutions are then used to inoculate sterile glass shell vials containing coverslips, as described above. The cells are then centrifuged and inoculated with virus or clinical samples, incubated, and fixed, as described above.

The viruses capable of infecting these cells are detected and identified using the methods described in Example 2 above. In these mixed cell cultures, the 292 cells are used to detect the presence of parainfluenza viruses and RSV, while the A549 cells are used to detect the presence of adenoviruses, and the mink lung cells are used to detect the presence of influenza viruses (*e.g.*, Influenza A and B).

EXAMPLE 7

Detection of HSV and *Chlamydia* in Mixed Cell Cultures

In this Example, mixed cell cultures are provided which allow the detection of two organisms commonly associated with sexually transmitted diseases. In these experiments, mink lung cells (*e.g.*, Mv1Lu) useful for the detection of HSV are mixed with McCoy cells useful for the detection of *C. trachomatis*.

First, McCoy cells and mink lung (*e.g.*, Mv1Lu) cells are grown in flasks, trypsinized, and mixed as described in previous Examples, such that a suitable dilution of mixed cells is produced. In preferred embodiments, the cells are diluted such that the mixed cells in culture will be in approximately the same proportions. These mixed cell dilutions are then used to inoculate sterile glass shell vials containing coverslips, as described above. The cells are then centrifuged and inoculated with samples (*e.g.*, clinical samples), incubated, and fixed, as described above.

The organisms capable of infecting these cells (*e.g.*, HSV infects the mink lung cells, while *C. trachomatis* infects the McCoy cells) are detected and identified using the methods described in Example 2 above. As with the other mixed cell culture systems, the presence of virus and/or *C. trachomatis* may be detected by other methods, such as the observation of CPE, animal inoculation, etc. Thus, it is not intended that the mixed cell culture assay systems of this Example or any of the preceding examples be limited to any particular method of microorganism detection, identification, and/or quantitation.

EXAMPLE 8
Evaluation of single cell cultures and mixed cell
cultures for detection of respiratory viruses

5 This Example evaluated different cell lines individually and in mixed cell culture.
The following cell lines were used in the exemplary shell vial with coverslip format: R-
mix (i.e., Mv1Lu and A549): C961023; Mv1Lu: C581023; A549: C561023; canine
kidney MDCK: C831022; NCI-H292: C591023; LLC-MK2: C861022; CV1:
C521023; pRHK: -CA-491016; MDCK/A549: C501022; MDCK/H292: C102303;
Mv1Lu-hF Clones numbers 15B, 17, 18, 29, 30, 35, 38 all 10-23-03.

10 The following reagents and virus strains were used: RM03T; Influenza A: WS,
Port Chalmers, Victoria, and Mai; Influenza B: Taiwan and G1; RSV: 031203 and
042403; Adenovirus: #1 and #5; Parainfluenza 1; Parainfluenza 2; Parainfluenza 3; D³
Kit: 091603; and Solution 1: 011303D.

15 Briefly, shell vials were all refed with 1 ml of RM03T. Virus dilutions were all in
RM03T. Shell vials were inoculated in duplicate with dilutions of each of the 7
respiratory viruses, i.e., influenza A, influenza B, RSV, adenovirus, parainfluenza 1,
parainfluenza 2, and parainfluenza 3. Shell vials were centrifuged for 1 hour at 700xg
then placed in a 35-37°C incubator. 24 hours, 48 hours and 72 hours post inoculation, a
set of shell vials were fixed and Stained with Solution 1 and D³.

20 The following is a key to the results shown in the following Tables 4-9: s = small.
B = Bursts. ~ = Approximately. TNTC = Too numerous to count. 1+ = 25% of
Monolayer infected. 2+ = 50% of Monolayer infected. 3+ = 75% of Monolayer infected.
4+ = 100% of Monolayer infected. N/A = Not Available. F = Field (there are 44 fields
per monolayer.)

Table 4
24 Hour Post Inoculation Results Using Influenza A Virus, Influenza B Virus,
RSV, and Adenovirus

Influenza A

		WS	Victoria	Port Chalmers	Mai
5	R-mix	99, 95	113, 127	197, 243	88, N/A
	MvILu	105, 114	142, 150	~5/F	169, 161
	A549	N/A	N/A	N/A	N/A
	MDCK	70 + ~8sB, 80 + 10sB	3sB + 71, 2sB + 69	2sB + 68, 2B + 88	2sB + 90, 3sB + 100
	H292	12, 5	17, N/A	12, 38	N/A, 7
10	LLC-MK2	11, 7	18, 19	38, 28	5, 2
	CV1	8, 7	23, 40	15, 12	N/A, 3
	pRHMK	~15B + ~10	5B + 50, 5B + 62	1+	5bigB, 1bigB + 5
	MDCK/A549	TNTC + Bursts	2B + 4sB + 75, 1sB + 104	145, 1B + 162	3bigB + 77, N/A
	MDCK/H292	TNTC + Bursts	67, 73	1sB + 84, 3B + 101	1bigB + 3sB + 69, 1B + 3sB + 101
15	15B	121, 124	3sB + 167, 6sB + 169	~300	~300
	17	168 + 2B, 150	132, 133	~200	5/F
	18	113, 120	3sB + 166, 171	241, 260	222, 200
	29	135, 152	109, 114	~200 + ~4sB	5/F
	30	109 + 1B, 125	133, 141	~300	6/F
20	35	75, 97	137, 140	~200	129, 170
	38	136, 132	113, 126	~300	5/F

Influenza B

		Taiwan	G1
25	R-mix	4/F	9/F
	MvILu	9/F	12/F
	A549	N/A	N/A
	MDCK	14B + 4/F, 8B + 4/F	9B + 6/F, 16B + 6/F
	H292	5, 8	13, N/A
	LLC-MK2	54, 50	16, 10
30	CV1	72, 80	45, N/A
	pRHMK	3B + 114, 2B + 128	39, 49
	MDCK/A549	2+	2+
	MDCK/H292	1+	1+
35	15B	11/F	10/F
	17	10/F	15/F
	18	12/F	12/F

29	9/F	9/F
30	10/F	9/F
35	7/F	8/F
38	9/F	10/F

5 **RSV**

	031203	042403
R-mix	54, 47	13, 16
MvILu	33, 27	3, 7
A549	34, 24	18, 23
MDCK	0, 0	0, 0
10 H292	23, 26	25, 22
LLC-MK2	30, 33	3, 6
CV1	20, 23	8, 9
pRHMK	0, 0	0, 0
MDCK/A549	23, 25	10, 13
15 MDCK/H292	28, 18	15, 19
15B	34, 38	N/A
17	43, 37	N/A
18	26, 30	N/A
29	18, 30	N/A
20 30	18, 24	N/A
35	21, 22	N/A
38	28, 40	N/A

Adenovirus:

	Adenovirus #1	Adenovirus #5
25 R-mix	20/F	~300
A549	20/F	216, 220
H292	5/F	46, 57
LLC-MK2	0, 0	0, 0
CV1	0, 5	0, 0
30 pRHMK	15/F	116, 160
MDCK	0, 0	0, 0
MDCK/A549	5/F	139, 124
MDCK/H292	55, N/A	4, 6

Table 5
24, 48 and 72 Hour Post Inoculation Results Using Parainfluenza 1

	24 hour	48 hour	72 hour
35 R-mix	64, N/A	143, 160	134, 143
MvILu	69, 73	118, 109	80, 88
A549	90, 194	111, 100	121, 110
MDCK	0, 0	0, 2	0, 0
40 H292	98, 111	158, 162	170, 159

	LLC-MK2	88, 106	163, 158	121, 117
	CV1	75, 66	68, 73	60, 72
	pRHKM	25sB + 25, 6sB + 40	4+	4+
	MDCK/A549	41, 50	62, 80	120, 122
5	MDCK/H292	38, 40	68, 75	68, 80
	15B	49, 54	110, 90	~100
	17	58, 63	119, 50B + 100	~120
	18	66, 62	87, 95	~100
	29	69, 65	63, 66	~70
10	30	23, 30	102, 115	~100
	35	47, 58	72, 75	~75
	38	50, 44	80, 85	~80

Table 6
24, 48 and 72 Hour Post Inoculation Results Using Parainfluenza 2

		24 hour	48 hour	72 hour
15	R-mix	66, N/A	1+	3+
	Mv1Lu	6, 10	~25	~20sB
	A459	210, 217	2+	4+
	MDCK	0, 0	0, 0	0, 0
	H292	116, 106	2+	4+
20	CV1	84, 94	1+	4+
	pRHKM	73, 80	2+	4+
	LLC-MK2	33, 29	~15B + 30, N/A	1+
	MDCK/A549	21, 28	~75	1+
	MDCK/H292	15, 24	~50	1+

Table 7
24, 48 and 72 Hour Post Inoculation Results Using Parainfluenza 3

		24 hour	48 hour	72 hour
	R-mix	5/F	TNTC	4+
	Mv1Lu	3/F	~50BB	~50BB
	A459	1+	4+	4+
30	MDCK	0, 0	~25	3/F
	H292	4/F	TNTC	4+
	CV1	1+	4+	4+
	pRHKM	1+	4+	4+
	LLC-MK2	4/F	TNTC	4+
35	MDCK/A549	~50	1+	4+
	MDCK/H292	~50	1+	4+

Table 8
48 Hour Post Inoculation Results Using Influenza A, Influenza B, RSV, and
Adenovirus

Influenza A:

		WS	Victoria	Port Chalmers	Mai
5	R-mix	79, 70	66, 72	128, 120	92, 67
	MvILu	60, 49	131, 127	94, 82	97, 94
	A549	N/A	N/A	N/A	N/A
	MDCK	4+	2+	2sB + 50, 47	1+
10	H292	8, 11	9, 13	2, 2	7, 10
	LLC-MK2	19, 25	1sB + 27, 31	55, 47	15, 10
	CV1	8, 21	24, 28	60, 48	6, 9
	pRHMK	3+	4+	4+	4+
15	MDCK/A549	4+	4+, 2+	1+	2+, 3+
	MDCK/H292	4+	1bigB, 2+	1B + 30, 2+	13B + ~100, 4+
	15B	77, 81	6sB + 75, 1+	140, 160	100, 119
	17	67, 65	76, 80	104, 113	124, 130
20	18	51, 61	66, 3sB + 100	133, 118	110, 105
	29	76, 60	2sB + 85, 91	143, 160	139, 115
	30	86, 70	56, 6B	150, 140	177, 160
	35	32, 40	43, 1sB + 52	87, 80	90, 83
	38	74, 1+	4sB + 81, 2sB + 88	4sB + 100, 6sB + ~100	118, 108

Influenza B:

		Taiwan	G1
25	R-mix	61, 64	120, 115
	MvILu	45, 65	120, 110
	A549	N/A	N/A
	MDCK	4+	4+
30	H292	1, 0	2, 2
	LLC-MK2	34, 33	11, 16
	CV1	23, 24	17, 19
	pRHMK	~10big B	35, 2big B
35	MDCK/A549	4+	4+
	MDCK/H292	3+	4+
	15B	100, N/A	147, 152
	17	80, 83	153, 149
	18	102, N/A	136, 141
	29	71, 73	74, 70
	30	83, 96	100, 94
	35	60, 53	108, 95
	38	70, 65	77, 72

RSV:

		031203	042403
	R-mix	60, 70	32, 39
	Mv1Lu	53, 50	2, 9
	A549	44, 43	23, 28
5	MDCK	0, 0	0, 0
	H292	40, 37	33, 54
	LLC-MK2	28, 35	13, 13
	CV1	26, 18	8, 7
	pRHMK	0, 0	0, 0
10	MDCK/A549	22, 6	12, 15
	MDCK/H292	19, 24	18, 18
	15B	39, 42	N/A
	17	68, 70	N/A
	18	54, 57	N/A
15	29	38, 50	N/A
	30	32, 31	N/A
	35	50, 32	N/A
	38	50, 66	N/A

Adenovirus:

		Adenovirus #1	Adenovirus #5
20	R-mix	3+	3+
	A549	3+	3+
	H292	3+	2+
	LLC-MK2	8/F	4/F
	CV1	70, 62	35, 40
25	pRHMK	2+	1+
	MDCK	2, 0	1, 0
	MDCK/A549	1+	1+
	MDCK/H292	1+	1+

Table 9
72 Hour Post Inoculation Results Using Influenza A, Influenza B,
RSV, and Adenovirus

Influenza A:

		WS	Victoria	Port Chalmers	Mai
	R-mix	<25	~25	~50	1sB + ~30
	Mv1Lu	<10	~25	~25	~25
35	A549	N/A	N/A	N/A	N/A
	MDCK	4+	2+, 4+	1+, ~50	3+, 2+
	H292	<5	<10	<5	<10
	LLC-MK2	<5	<25	<5	<10

	CV1	<5	<25	~10	<10
	pRHMK	4+	4+	4+	4+
	MDCK/A549	4+	1+, 4+	4+	4+
	MDCK/H292	4+	~20B, 4+	1+, 4+	4+
5	15B	~25	12sB, ~25	~25	~25
	17	~25	<5	~25	~25
	18	<10	3B + 50, ~25	<25	~50
	29	~10	~15sB, ~25	<25	~25
	30	<10	<10	<25	~50
10	35	<10	<25	<5	~25
	38	~2sB + ~50	50, 25	~25	~50

Influenza B:

		Taiwan	G1
	R-mix	8, 17	~30
	MvILu	0, 0	<5
15	A549	N/A	N/A
	MDCK	4+	4+
	H292	<5	<5
	LLC-MK2	7, 10	5, 7
	CV1	<5	<5
20	pRHMK	4+	<5, 4+
	MDCK/A549	4+	4+
	MDCK/H292	4+	4+
	15B	0, 0	<5
	17	<5	<5
25	18	0, 0	<5
	29	0, <5	<5
	30	0, <5	<5
	35	<5	<5
	38	<5	<5

RSV:

		031203	042403
	R-mix	59, 50	24, 27
	MvILu	45, 52	7, 20
	A549	60, 56	24, 36
	MDCK	0, 0	0, 0
35	H292	26, 30	50, N/A
	LLC-MK2	31, 28	8, N/A
	CV1	25, 31	13, 19
	pRHMK	2, 1	3, 5
	MDCK/A549	13, 16	17, 19
40	MDCK/H292	21, 30	12, 16
	15B	39, 45	N/A
	17	38, 42	N/A

18	40, 47	N/A
29	30, 35	N/A
30	34, 30	N/A
35	39, 35	N/A
38	40, 43	N/A

Adenovirus:

	Adenovirus #1	Adenovirus #5
R-mix	3+	4+
A549	4+	4+
H292	2+	2+
LLC-MK2	1+	1+
CV1	4/F	4/F
pRHMK	2+	2+
MDCK	0, 0	0, 0
MDCK/A549	2+	2+
MDCK/H292	1+	1+

The above data show that the mixed cell cultures of MDCK+A549 and MDCK+H292 showed comparable sensitivity to R-mix, i.e., Mv1Lu and A549 cells with respect to detecting the seven exemplary respiratory viruses: respiratory syncytial virus (RSV), adenovirus, parainfluenza 1 virus, parainfluenza 2 virus and parainfluenza 3 virus. In one embodiment, mixtures of MDCK with one or more of A549 and H292 cells may preferably be used at 24 hours in culture since, by 48 and 72 hours, the MDCK almost completely outgrew the other cell lines.

EXAMPLE 9

Comparison of MDCK and Mv1Lu cells inoculated with Influenza A and B

This example was carried out to determine the ability of MDCK and Mv1Lu cells to propagate strains of Influenza A and B. Cultures were tested using duplicate monolayers at 24, 48 and 72 hours post inoculation. Where virus is replicating, more positive cells (such as those detected by fluorescence) were expected by the inventors to be observed at the 48 and 72 hour time points compared to the zero time point of inoculation.

The following exemplary cells and viruses were used: MDCK lot C830807; Mv1Lu lot C580807R; RM03T lot 070903E; ELVIS Solution 1 lot 061203 (Diagnostic Hybrids, Inc., Ohio, USA); Influenza A and Influenza B components from D³ Kit lot 011303; ELVIS Mounting Fluid lot 011603A (Diagnostic Hybrids, Inc., Ohio, USA).

Briefly, cell cultures of MDCK and Mv1Lu shell vials with coverslips were used. All cultures were refed with 1ml of RM03T. Virus stocks were rapidly thawed in a 35-37°C bath and diluted to a working stock in RM03T. Each culture was inoculated in duplicate with 200µl of each working virus stock. All cultures were centrifuged at 700xg for 1 hour. All cultures were placed in a 35-37°C incubator. A set of each was processed according to the D³ Kit product insert at 24, 48 and 72 hours post inoculation.

Table 10
Comparison of MDCK and Mv1Lu cells Using Influenza A and Influenza B

	Virus/strain/lot #	MDCK			Mv1Lu		
		Day 1	Day 2	Day 3	Day 1	Day 2	Day 3
10	Flu A: Denver: 112701N	156, 129 +1B	~15/F + 5BB	4+ CPE	<i>179, 164</i>	<i>110, 115</i>	<i>~25</i>
	Flu A: Aichi: 112701K	<i>189, 206 + 1B</i>	<i>100, 80</i>	<i>~100 + 1B</i>	<i>~7/F</i>	<i>~6/F</i>	<i>~60</i>
15	Flu A: PR: 111201D	114 + 5B, 118 + 4B	~1+CPE	<i>~50</i>	<i>147, 158</i>	<i>50, 42</i>	<i>~4</i>
	Flu A: Victoria: 121800	121 + 2B, 106	1+ CPE	<i>~30 + ~5B</i>	171, 208	1+ CPE	1+ CPE
	Flu A: WS: 111201E	118 + 1BB, 120+ 7B	3+ CPE	4+ CPE	87, 122	1+ CPE	3+ CPE
20	Flu A: Portchalmers: 112701	59, 68	~5 + 1B	~100 + ~3B	<i>105, 98</i>	<i>~60</i>	<i>~100 + ~2B</i>
	Flu A: Mal: 112701L	106 + 1B, 118 +3B	~50 + ~6B	4+ CPE	<i>176, 175</i>	<i>~100</i>	<i>~100</i>
25	Flu A: HongKong: 112701M	112 + 1B, 85 + 1B	~50 + 2B	<i>~100</i>	<i>160, 170</i>	<i>~100 + ~10B</i>	<i>~100</i>
	Flu A: NJ: 102699	134 + 2B, 113 + 3B	2+ CPE	4+ CPE	<i>225, 190</i>	<i>~85</i>	<i>~75</i>
30	Flu B: GL: 112701S	~5/F + 1B	3+ CPE	4+ CPE	<i>~10/F</i>	<i>~50</i>	<i>~50</i>
	Flu B: Taiwan: 112701R	~5/F + ~1B/F	3+ CPE	4+ CPE	<i>~8/F</i>	<i>~50</i>	<i>~10</i>
35	Flu B: HongKong: 020402B	81, 82	~10	~100	<i>125, 140</i>	<i>~10</i>	<i>~40</i>
	Flu B: Mass.: 112701Q	<i>52, 60</i>	<i>~20</i>	<i>~20</i>	<i>199, 216</i>	<i>~30</i>	<i>~20</i>
40	Flu B: Maryland: 112701P	~75B + tntc S	3+ CPE	4+ CPE	<i>~20/F</i>	<i>~50</i>	<i>~100 + 1B</i>

Flu B: Russia: 112701FF	~8/F + ~20B	2+ CPE	4+ CPE	~10/F	~40	~20
----------------------------	----------------	--------	--------	-------	-----	-----

123 = number of single fluorescent cells.

B = Burst of fluorescent cells. Usually 100 or more together.

BB = Big Burst. Usually described by percentage of monolayer covered.

S = Single cells.

~ = approximately. Usually used as an average of both monolayers.

+ = and. Unless used before CPE. (See CPE below).

5/F = 5 single cells per field. There are 44 fields per coverslip.

tntc = Too numerous to count.

CPE = cytopathic effect. This ranges from 1+ to 4+ with 1=25%, 2=50%, 3=75% and 4=100% of cells infected.

Bold = increasing titer. (virus replication)

Italic = decreasing titer. (no virus replication)

In the above experiments, 11/15 virus strains were propagated in the MDCK cell line. Influenza A: Aichi and Flu B: Mass. had lower titers on days 2 and 3. Influenza A Hong Kong and Influenza A Port Chalmers did not have any significant change in virus titer from 1 to 3 days of culture. The data shows that 2/15 virus strains were propagated in the Mv1Lu cell line. They were Influenza A: Victoria and Influenza A: WS. 11/15 virus stocks cultured in the Mv1Lu lost titer after 24 hours. 2 virus strains remained the same titer over the 3 days in the Mv1Lu cell line. The day 1 results showed the Mv1Lu cells to be slightly more sensitive than MDCKs as measured by the number of positive individual cells, however, the MDCKs were the only cell line to show bursting at 24 hours. Based on this data, there is no significant difference on day 1 initial titer between the Mv1Lu and MDCK cell lines. Surprisingly, MDCK cells detect and produce influenza A and B at higher levels than the Mv1Lu cells.

Thus, the use of MDCK in single cell culture and in mixed cell culture with one or more of H292 and A549 is useful for identifying low levels of influenza A virus and influenza B virus at the exemplary times of 48 and 72 hours post-inoculation, as well as for producing influenza A virus and influenza B virus.

EXAMPLE 10

Materials And Methods

The following is a brief description of the exemplary materials and methods used in the subsequent Examples.

A. Virus

A seed stock of SARS-CoV Urbani that was passaged twice in Vero E6 cells provided by the Centers for Disease Control and Prevention, Atlanta, GA. This virus was amplified by two passages in Vero E6 cells to establish a high titer stock (passage 4) that was utilized for all experiments. SARS-CoV was titered in Vero E6 cells by TCID₅₀. Briefly, cells were plated in 96-well plates (Falcon, Becton Dickson) at a density of 4×10^5 cells/well in 150 μ l of medium. Virus was serially diluted by half logs from 10^0 – 10^{-7} in culture medium containing 2% antibiotic-antimycotic (Invitrogen Corporation, Carlsbad, CA). 100 μ l of each dilution was added per well and cells were incubated 3-4 days at 37°C.

B. Cell lines

The following Table lists exemplary cell lines that were used and/or equivalent cells that may be used in the invention's methods, and that are publically available (*e.g.*, from the American Type Culture Collection (ATCC), Rockville, Maryland, and Diagnostic Hybrids, Inc. (DHI), Athens, Ohio; Cell Bank, Ministry of Health and Welfare, Japan):

Table 11
Exemplary Cells Useful In The Invention

Cells	Source
Vero E6	ATCC # CRL-1586 DHI # 67-0102
MRC-5	ATCC # CCL-171 DHI # 51-0102
BHK-21	ATCC # CCL-10 DHI # 89-0102
MDCK	ATCC # CCL-34 DHI # 83-0102
HRT-18 (HCT-18)	ATCC # CCL-244
Mv1Lu	ATCC # CCL-64 DHI # 58-0102

	CMT-93	ATCC # CCL-223
	AK-D	ATCC # CCL-150
	A549	ATCC # CCL-185
		DHI # 56-0102
	HEL	DHI # 88-0102
5	pRHMK	DHI # 49-T025
		DHI # 49-0102
	pCMK	DHI # 47-T025
		DHI # 47-0102
	L2	ATCC # CCL-149
	R-Mix	DHI # 96-T025
	HEK-293T	ATCC # CRL-1573;
		CRL- 11264, CRL-
		11270; Pear, et al., PNAS
		USA, Vol 90, pp
		8392-8396 Sept. 1993;
		DuBridge et. al., Mol.
		Cell. Biol. Vol 7, pp
		379-387, 1987;
		University Dr. Yoshi
		Kawaoka, Univ.
		Wisconsin, Madison.
10	Huh-7 (JTC-39)	CellBank #JCRB0403

15 R-Mix (R-Mix FreshCells™, Diagnostic Hybrids, Inc., Ohio) is a mixed monolayer of mink lung cells (strain Mv1Lu) and human Adenocarcinoma cells (strain A549). the hAPN expression construct used to create BHK21/hAPN and CMT-93/hAPN was previously described (Wentworth, et al., 2001). Further description of Huh-7 cells is in Nakabayashi et al., Cancer Res., 42: 3858-3863, 1982; Nakabayashi et al., Gann, 75: 151-158, 1984; and Nakabayashi et al., Cancer Res., 45:6379-6383, 1985.

Vero E6, 293T, L2, AK-D, A549, pCMK, pRhMK, Mv1Lu, CMT-93, and R-mix were maintained in Dulbecco's modified Eagle Medium (DMEM) (Invitrogen Corp.) supplemented with 10% fetal bovine serum (FBS) (Hyclone, Logan, UT) and

2% antibiotic-antimycotic. MDCK cells were maintained in DMEM high glucose (Invitrogen Corp.) supplemented with 5% FBS and 2% antibiotic-antimycotic. HEL cells were maintained in Modified Eagle's Medium (MEM) supplemented with 10% FBS and 2% antibiotic-antimycotic. HRT-18 cells were maintained in RPMI 1640 (Invitrogen Corp.) supplemented with 10% horse serum (Hyclone), 1 mM MEM sodium pyruvate (Invitrogen Corp.) and 2% antibiotic-antimycotic. Huh-7 cells were maintained in DMEM supplemented with 20% FBS and 2% antibiotic-antimycotic. MRC-5 cells were maintained in MEM supplemented with 10% FBS, 1 mM sodium pyruvate, 0.1 mM MEM nonessential amino acids (Invitrogen Corp.) and 2% antibiotic-antimycotic. BHK-21 cells were maintained in DMEM supplemented with 10% FBS and 5% tris phosphate buffer (Invitrogen Corp.).

C. PCR assay

G3PDH, genomic SARS-CoV RNA (gRNA) and subgenomic RNA (sgRNA) were detected using multiplex one-step RT-PCR. Oligonucleotide primers used to amplify the different targets were as follows: G3P-279 (sense) 5' CATCACCATCTTCCAGGAGC-3' (SEQ ID NO:7) binds at nt 279-299; G3P-1069R (antisense) 5'- CTTACTCCTTGGAGGCCATG -3' (SEQ ID NO:8) binds at nt 1069-1049; SARS-21,263 (sense) 5'- TGCTAACTACATTTTCTGGAGG-3' (SEQ ID NO:9) binds at nt 21,263-21,284 of SARS-Urbani; SARS-21,593R (antisense) 5'-AGTATGTTGAGTGTAATTAGGAG-3' (SEQ ID NO:10) binds at nt 21,593-21,571 of SARS-Urbani; and SARS-1 (sense) 5'- ATATTAGGTTTTTACCTACCCAGG-3' (SEQ ID NO:11) binds at nt 1-24 of SARS-Urbani. Amplification was carried out using the Qiagen® OneStep RT-PCR kit (Qiagen) according to the manufacturer's protocol. Briefly, each reaction consisted of 2 µg of total RNA isolated using TRIZOL® Reagent (Invitrogen), 400 µM dNTPs, 200 nM of each G3PDH primer, 400 nM SARS-1, 400 nM SARS-21,263, 600 nM SARS-21,593R and 2 µl Qiagen enzyme mix. The cycling parameters were: 50°C for 30 min, 95°C for 15 min, 35 cycles of 94°C for 30 s, 57-58°C for 30 s, 72°C for 1 min, followed by 10 min at 72°C in an Eppendorf Mastercycler gradient (eppendorf). Amplification products were analyzed by electrophoresis through a 1.5% agarose gel and visualized by ethidium bromide staining. All primers were synthesized by the Molecular Genetics Core (David Axelrod Institute, Wadsworth Center, Albany, NY).

D. Cell infection

Cells seeded at a density of 2×10^6 in T25 flasks (Falcon, Becton Dickson) were inoculated with virus at an MOI of 0.001 in a final volume of 1 ml and were incubated 1 h at 37°C. Virus was removed and 5 ml fresh medium added to each flask. Cells were maintained at 37°C throughout the experiment. At 1, 24 and 48 h post-inoculation (p.i.), cells were observed for CPE, supernatants were collected for subsequent titration and total RNA was extracted using TRIZOL[®] Reagent (Invitrogen Corp.). RNA was quantitated by spectrophotometer (Eppendorf).

EXAMPLE 11

Exemplary multiplex RT-PCR assay for the detection of SARS-CoV replication

A RT-PCR assay for the detection of SARS-CoV replication was developed. Replication of corona- and arteri- virus RNA occurs through discontinuous synthesis, thought to occur during negative strand synthesis, generating 3' co-terminal nested subgenomic RNAs (sgRNA). The inventors identified targets within the genome for amplification. Oligonucleotide RT-PCR primers were designed that amplify genomic SARS-CoV RNA (gRNA) or the sgRNA that is specific to the leader-body junction. Because genomic RNA is present in input virus, the inventors probed for sgRNA, which is indicative of virus entry and/or replication initiation. Genomic RNA was detected by amplifying a region between the 1b coding region of the polymerase gene and the sequence encoding the Spike (S) glycoprotein. Subgenomic RNA was detected using a primer specific to the leader sequence in conjunction with the reverse primer in S that was used for the gRNA detection. G3PDH primers, designed to amplify G3PDH from multiple species, served as a positive control for RNA integrity and cDNA production.

To evaluate the RT-PCR assay, Vero E6 cells were inoculated with serial dilutions of SARS-CoV ranging from an MOI of 10^0 to 10^{-8} TCID₅₀/cell. Total RNA was extracted at 1 and 24 h post-inoculation (p.i.). At 1 h p.i. gRNA was detected in cells inoculated with virus at an MOI of 10^0 to 10^{-2} , as indicated by a band at 300bp (Fig. 1). Subgenomic RNA was not detected (180 bp). However, at 24 h p.i. both gRNA and sgRNA, 300 bp and 180 bp respectively, were detected in cells inoculated with an MOI of 10^0 to 10^{-5} . The sgRNA amplicon was confirmed to correspond to the S leader-body junction sgRNA by sequence analysis (Thiel, et al., 2003, J.Gen.Virol. 84:2305-2315). Genomic RNA was visible at 24 h p.i. in cells inoculated with an MOI of 10^{-7} , however this was not seen in repeated experiments. The decrease in

amplified G3PDH (~800bp) as seen in lanes 1-6 at 24 h p.i. was consistent between repeated experiments. The decrease in G3PDH may be a result of the RT-PCR conditions, which were optimized to favor amplification of SARS-CoV gRNA and sgRNA. Individual amplicon were amplified by PCR of cDNAs from the same samples and G3PDH was consistently detected. Additionally, the decrease in G3PDH may be due to cell death, which is seen in Vero E6 cells. G3PDH was included as a control for template concentration and RNA integrity, and was always detected in the absence of viral RNA.

This data demonstrates that the exemplary multiplex RT-PCR assays is sensitive for detection of SARS-CoV infection.

EXAMPLE 12

Cytopathic Effect Does Not Always Correlate With SARS-CoV infection

This example shows replication of SARS-CoV, as detected by sgRNA and virus titers, in the absence of CPE. In particular, significant CPE was not observed in pRhMK, pCMK, R-mix (Mv1Lu and A549), Mv1Lu, HEK-293T, and Huh-7 cells at 5 days post infection, although virus titers as well as SARS-CoV sgRNA were actually increased within 24 hours post infection (Table 12).

TABLE 12
Susceptibility of cells to SARS-Coronavirus

		SARS-CoV			
	Cell	Species of Origin	sgRNA	CPE	Titer ^a
5	VeroE6	African green monkey	+	+	2.4x10 ⁷
	pRhMK	Rhesus macaque	+	-	5.6x10 ⁵
	PCMK	Cynomolgous macaque	+	-	7.8x10 ⁴
	R-Mix	Mink and Human	+	-	7.8x10 ³
	A549	Human	-	-	<1 ^b
10	Mv1Lu	Mink	+	-	2.5x10 ⁴
	HEL	Human	-	-	<1
	MRC-5	Human	-	-	<1
	MDCK	Canine	-	-	<1
	AK-D	Feline	-	-	ND ^c
15	L2	Murine	-	-	ND
	HRT-18	Human	-	-	ND
	CEF	Chicken	-	-	ND
	HEK-293T	Human	+	-	5.6x10 ³
	Huh-7	Human	+	-	1.3x10 ⁵
20	CMT-93	Murine	-	-	ND
	CMT-93/hAPN	Murine	-	-	ND
	BHK-21	Syrian hamster	-	-	<1
	BHK-21/hAPN	Syrian hamster	-	-	ND

25 ^a Titer = TCID₅₀/ml at 48hr post-inoculation.

^b Below the limit of detection

^c Titer not determined.

From the above, it is clear that the present invention provides many advantages over presently used methods in diagnostic microbiology.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in diagnostic microbiology and virology, cell culture, and/or related fields are intended to be within the scope of the following claims.